



Research article

**Antimicrobial and antioxidant studies of Schiff base, 2-(2-hydroxy-2-methyl-1-phenylpropylidene) hydrazine carbothioamide and its mixed ligand cd (II) complexes**Vanitha Sanambatl<sup>1</sup>, Sathish Kumar Nara<sup>2</sup>, Balaji Hari<sup>2</sup>, Varada Reddy Ammireddy<sup>2</sup>, Saritha Nimmakayala<sup>3\*</sup><sup>1</sup>J. N. T. U. Anantapur, ananthapuramu, India<sup>2</sup>S.V. University, Tirupati, Andhra Pradesh, India<sup>3</sup>J. N. T. U. A. College of Engineering, Kalikiri, Andhra Pradesh, India**ABSTRACT**

New mononuclear mixed ligand Cadmium(II) complexes of the type  $[Cd(L)(diimine)]$  (1,2) [where L = 2-(2-hydroxy-2-methyl-1-phenylpropylidene)hydrazinecarbothioamide; diimine = 2,2'-bipyridine (1), 1,10-phenanthroline (2)] have been synthesized and characterized by spectroscopic techniques such as FT-IR, UV-Visible, and <sup>1</sup>H and <sup>13</sup>CNMR Spectroscopy. From the investigations of spectral data, it is evident that the heterocyclic bases (2,2'-bipyridine and 1,10-Phenanthroline) act as neutral bidentate ligand coordinating to the metal ion through two nitrogen donor atoms addition to azomethane nitrogen, thiolate Sulphur, and hydroxyl oxygen in the Schiff base ligand. The synthesized Schiff base metal chelates have been screened for their anti-microbial activities using the agar well diffusion method against different selected types of bacteria and fungi in addition to antioxidant activity. The prepared Schiff base ligand and its metal complexes exhibited good antimicrobial and antioxidant activities. The antibacterial and antifungal efficacy of Complex-1 was higher than that of all the prepared compounds. In the case of antioxidant activity, Complex-2 has stronger scavenging activity among them.

**Keywords:** Cd (II) metal complexes, 2,2'-Bipyridine, 1,10-Phenanthroline, Antimicrobial activities.

Received - 20-11-2021, Accepted- 20-01-2022

**Correspondence:** Dr. N Saritha Nimmakayala\* ✉ [sarithanphd@gmail.com](mailto:sarithanphd@gmail.com)

Department of Chemistry, J. N. T. U. A, College of Engineering, Kalikiri, Andhra Pradesh, India.

**INTRODUCTION**

Because of the pharmacological properties of both the ligands and the complexes, the synthesis of transition metal complexes with thiosemicarbazone ligands has received a lot of attention [1]. Thiosemicarbazone ligands typically coordinate to metals via oxygen, nitrogen and Sulphur donor atoms in their (N, S) bidentate form or (N, N, S or O, N, S) tridentate form, resulting in metal complexes with varying molecular geometries [2]. Such complexes are especially important because of their potential biological benefits, such as anticancer [3], fungicidal [4], antibacterial [5], antiviral [6], antifungal [7,8], antitumor [9], and other biological activities [10], particularly with the first row of transition metal complexes. Free ions are more hazardous than their metal complexes due to the lack of bioavailability of the metal. Taking this into account, researchers have been looking at the biological functions of new cadmium Schiff base metal complexes [11-16]. These have also been proved as effective antibacterial agents due to the presence of active biological strains [17]. With this perspective, the current research focused on cadmium based mixed ligand metal complexes.

Because of their structural diversity and wide range of applications, mixed-ligand complexes receive a lot of attention in coordination chemistry. Mixed-ligand compounds also have a wide range of biological applications [18].

In the present work, 2-(2-hydroxy-2-methyl-1-phenylpropylidene) hydrazine carbothioamide and heterocyclic bases like 2, 2'-bipyridine and 1,10-phenanthroline have been used as co-ligands for the synthesis of cadmium (II) complexes was carried out. Synthesis, structures, spectroscopy and antimicrobial aspects of complexes are investigated.

**MATERIALS AND METHOD****Chemicals**

The starting materials such as 2-Hydroxy-2-methylpropiophenone, thiosemicarbazides and heterocyclic bases like 2,2'-bipyridine, and 1,10-Phenanthroline were acquired from Sigma Aldrich, India. Cd (II) salt in the form of CdCl<sub>2</sub>·2H<sub>2</sub>O was purchased from S. D. Fine chemicals. The organic solvents like acetonitrile, methanol, ethanol, dichloromethane was purchased from Merck, which were not subjected to any further purification process.

### Synthesis of the ligand and their Cd (II) complexes

Synthesis of Schiff base ligand, 2-(2-hydroxy-2-methyl-1-phenylpropylidene) hydrazine carbothioamide

The Schiff base ligand was synthesized as per the reported literature protocol [19,20]. To the solution of 2-hydroxy-2-methyl propiophenone (0.01 mol) in methanol, thiosemicarbazone (0.01 mol) was added slowly. The above mixture was refluxed for 3 hours with constant stirring at a temperature of 70°C. Then the reaction mixture was evaporated at room temperature. Colorless compound was separated out. It was washed with methanol. Purity was checked by Thin Layer Chromatography (TLC).

Synthesis of the mixed ligand Cadmium (II) Complexes (1,2)

Schiff base ligand (L) (0.001 mol) was dissolved in 20 ml of

methanol followed by methanolic CdCl<sub>2</sub> (0.001 mol) was added. The mixture was refluxed for 1 h. with constant stirring. The yellow-colored solution was formed. To the resultant reaction mixture, heterocyclic base (2,2'-bipyridine) (0.001 mol) which is dissolved in a mixture of dichloromethane and methanol was added slowly. The mixture was refluxed for 6 hours with constant stirring. The lemon-yellow colored precipitate was formed after that time period which was then filtered, and evaporated using Rota evaporator. The yellow color precipitate was obtained. The precipitate was washed several times with cold ethanol and dried under vacuum pump over anhydrous CaCl<sub>2</sub>. The schematic representation for synthesis of Schiff base ligand L and Cd (II) metal complexes are presented in figure 1 and figure 2. The same procedure was followed for the Schiff base mixed ligand complex with 1,10-Phenanthroline.

Figure 1: Schematic preparation of Schiff base ligand





Figure 2: Schematic preparation of mixed ligand Cd (II) metal complexes



### ANTIMICROBIAL ASSAYS

#### Preparation of media

The antibacterial activity of ligand and their metal chelates is assessed using the agar well diffusion method. These assays are used to create a dosage response curve for a specific bacterial strain

versus a specific test chemical. Two Gram positive bacteria strains, Bacillus subtilis and Staphylococcus aureus and two Gram negative bacteria strains, Escherichia coli and Klebsiella pneumoniae are tested for antibacterial activity. Three fungal strains, Aspergillus

Niger, *Aspergillus flavus*, and *Rhizopus oryzae* were used to test antifungal activity. Antibacterial and antifungal reference medications were Streptomycin and Fluconazole, respectively.

#### Antioxidant activity

The 2,2-diphenyl-1-picryl-hydrazyl-hydrate (DPPH) free radical technique was used to examine the antioxidant and free radical scavenging properties of the produced compounds in vitro. In methanol, a 0.1 M DPPH solution was prepared. The crude extract was then dissolved in 10% Dimethylsulfoxide (DMSO) at various concentrations (20 - 100 g mL<sup>-1</sup>) and 500 µL of the DPPH solution was combined with 1 mL of the crude extract. These solutions were mixed together and incubated in the dark for about 30 minutes at room temperature. At 517 nm, the absorbance of ascorbic acid was


compared to that of a blank with no scavenger activity.

## RESULTS AND DISCUSSION

### <sup>1</sup>H NMR spectra

The <sup>1</sup>H NMR spectra (Model: Bruker 400) of the prepared Schiff base was recorded using TMS as an internal standard in DMSO-d<sub>6</sub> as a solvent for ligand. This was shown in figure 3. The peak around 1.0 ppm can assign to the six methyl hydrogens in the compound. The alcoholic hydrogen was seen at 5.2 ppm. The peaks from 7.2 ppm to 7.7 ppm can assign to aromatic hydrogens. The -NH group and -NH<sub>2</sub> group hydrogen atoms of the semi thiocarbazide fragment were seen at 8.0 ppm and 8.3 ppm respectively. The <sup>1</sup>H NMR spectra of compound clearly insights into formation of the imine group between ketone and amine to give Schiff base ligand.

Figure 3: <sup>1</sup>H NMR Spectra of Schiff base




### <sup>13</sup>C NMR spectra

The <sup>13</sup>C NMR spectra (Model: Bruker 400) of the prepared Schiff base was recorded using TMS as an internal standard in DMSO-d<sub>6</sub> as a solvent for ligand. The spectral image was shown in figure 4. <sup>13</sup>C NMR spectra is an important tool to elucidate the structure of the prepared Schiff base ligand. The two methyl carbons attached to tertiary carbon atom was seen at 28 ppm. The tertiary

carbon that adjacent to imine group was seen at 72 ppm. The aromatic carbon atoms are seen in the range of 128 ppm to 134 ppm. The most importantly, the peak at 158 ppm was given to imine carbon atom. The -C=S group carbon atom was assigned to the peak at 183 ppm. From the Carbon NMR spectra, it was concluded that the formation of Schiff base complex was successful using the starting ketone and amine.

Figure 4: <sup>13</sup>C NMR Spectra of Schiff base



### FT-IR analysis

Infrared spectra's were recorded using Bruker alpha spectrometer in the wave number range of 4000 – 400  $\text{cm}^{-1}$ . In the uncoordinated ligand, an important strong band appears at 1666  $\text{cm}^{-1}$  attributing to free azomethine group which confirms the formation of Schiff base by condensation of ketone and amine, but in metal complexes a negative shift up to 1658  $\text{cm}^{-1}$  suggests coordination of the imine nitrogen to metal centers. This may occur due to decrease in bond strength of imine bond and simultaneous increase in bond strength between azomethine nitrogen and metal center. The  $-\text{NH}_2$  stretching frequency was clearly seen around 3200  $\text{cm}^{-1}$  to 3300  $\text{cm}^{-1}$  and sometimes coupled with  $-\text{OH}$  and  $-\text{NH}$  stretching's. The aliphatic  $-\text{C}-\text{H}$  stretching was seen around 2900  $\text{cm}^{-1}$ . The aromatic  $-\text{C}=\text{C}-$  stretching was seen at 1490  $\text{cm}^{-1}$ . The  $-\text{C}=\text{S}$  stretching was seen at 1389  $\text{cm}^{-1}$ . In the case of metal complexes, the metal complexes show absorption peaks in the region 420 - 427  $\text{cm}^{-1}$  corresponding to M-N and 655 - 658  $\text{cm}^{-1}$  for M-O vibrations confirming the bond formation between azomethine nitrogen, hydroxyl oxygen and metal ion. Another absorption band in the range of 3300 - 3550  $\text{cm}^{-1}$  in complexes marks the presence of coordinated or lattice water. This is presented in Table 1 and the FT-IR Spectra of Schiff base ligand is shown in figure 5.

Figure 5: FT-IR Spectrum of Schiff base ligand




Table 1: FT-IR Spectral data for Ligand and its Cd (II) Metal Complexes


Compound	Wavenumber ( $\text{cm}^{-1}$ )				Bands due to heterocyclic bases
	$\nu(\text{C}=\text{N})$	$\nu(\text{N}-\text{N})$	$\nu(\text{C}=\text{S})$	$\nu(\text{Cd}-\text{N})$	
L	1666	1099	1389, 871	--	--
1	1658	1089	1375, 863	427	1499, 657
2	1656	1088	1379, 864	425	1498, 656

### Electronic spectra

UV-Visible spectra were recorded in the range of 200 – 800 nm on Cary 5000 version spectrophotometer at room temperature using solutions in Dimethylformamide (DMF). The electronic absorption spectra of Cd (II) complexes recorded in DMF solution is shown in figure 6. The bands observed indicate  $\pi \rightarrow \pi^*$  transitions,

confirming metal center binding with the Schiff base, 1,10-phenanthroline / 2,2'-bipyridine. The ligand's electronic spectrum shows bands at 274 nm and 347 nm that corresponds to  $\pi \rightarrow \pi^*$  transitions and  $n \rightarrow \pi^*$  transitions of the azomethine and thioamide functions, respectively. Because of the coordination of the ligand with metal ions, these transitions were shifted to higher or lower frequencies [21]. The shift of the  $\pi \rightarrow \pi^*$  bands to longer wavelength regions in complexes is caused by the weakening of the  $\text{C}=\text{S}$  bond and the enhancement of the conjugation system during complexation [22]. There were no detectable absorptions above 500 nm in DMF solution, indicating the absence of d-d bands, which is consistent with the  $d^{10}$  configuration of the Cd (II) ion.

Figure 6: Electronic Spectrum of ligand and its complexes (1,2)



### Biological activity


The antibacterial activity of synthesized compounds was tested in-vitro using the agar-well diffusion method against a variety of harmful bacteria and fungus strains.

#### Antibacterial Activity

For both bacteria, the ligand antibacterial activity was shown to be weaker than that of the standard reference antibiotic (Streptomycin), however the Cd (II) complexes were found to be more effective and reached good activity when compared to the standard drug. This indicates, the formation of complexes with ligands and metal ions was happened. The antibacterial activity of complex-1 was higher than that of remaining complex and the Schiff base ligand. Because of the chelation interaction between Schiff bases and ligand complexes, complex-1 demonstrated good activity for both positive and negative bacteria and is shown in Table 2 and figure 7. Lipophilicity, which controls the rate at which molecules enter the cell, is similarly affected by coordination.

Table 2: Antibacterial activity of compounds [Zone of inhibition (mm)]

Compounds	Gram's (+)		Gram's (-)	
	Bacillus subtilis	Staphylococcus aureus	Escherichia coli	Klebsiella pneumoniae
L	0.8	0.8	0	0.8
1	1.5	1.2	1.2	1.0
2	0.9	0.9	1.1	1.0
Standard Drug (Streptomycin)	1.7	2.8	3.4	2.3


**Figure 7:** Zone of inhibition values for antibacterial activity**Antifungal activity**

Fluconazole was considered as the standard drug for evaluating antifungal activities of the ligand and its metal complexes. Complex-1 had moderate antifungal efficacy against three different fungal strains (*Aspergillus Niger*, *Aspergillus flavus*, *Rhizopus oryzae*). Complex-2, on the other hand, shows lesser efficacy against *Aspergillus flavus* and *Rhizopus oryzae*. Complex-2 is not reactive for *Aspergillus Niger*. Henceforth, Complex-1 shown better antifungal efficacy. Table 3 and Figure 8 depicts inhibition

zone values of the complexes and the ligand in comparison with the standard drug (Fluconazole).

**Table 3:** Antifungal activity of compounds [Zone of inhibition (mm): Paper Disc Method]


Compounds	<i>Aspergillus Niger</i>	<i>Aspergillus flavus</i>	<i>Rhizopus oryzae</i>
L	0.7	0.6	0.8
1	0.9	1.1	1.5
2	0	1.2	1.1
Standard Drug (Fluconazole)	0.7	2.1	2.1

**Figure 8:** Zone of inhibition values for antifungal activity**Antioxidant activity**

Using the DPPH assays method, the ability of the newly synthesized Schiff bases ligands and their complexes to scavenge free radicals may be assessed, and the findings are shown in Figure 9. Because of its stability and simplicity, the DPPH radical is frequently used to evaluate antioxidant activity quickly. The creation of stable 1,1-diphenyl-2-picrylhydrazine resulted in a decrease in the

intensity of the DPPH band, demonstrating the ability of metal complexes and Ascorbic acid (used as a standard) to scavenge free radicals [23]. Increased antioxidant activity is indicated by a decrease in absorbance and a decrease in the IC<sub>50</sub> value [24,25]. The IC<sub>50</sub> values of the test compounds were observed to be in the order L > 1 > 2, indicating that complex-2 has a stronger scavenging activity, while complex 1 has a moderate scavenging activity.



**Figure 9:** IC<sub>50</sub> values of Schiff base ligand and Cd (II) metal complexes (1,2)

## CONCLUSION

In this paper, reported the synthesis of two Schiff base mixed ligand Cd (II) metal complexes namely [Cd(L)(diimine)] (1, 2) [where L = 2-(2-hydroxy-2-methyl-1-phenylpropylidene) hydrazine carbothioamide; diimine = 2,2'-bipyridine (1), 1,10-phenanthroline (2)]. The complexes were elucidated by the use of various techniques. The antibacterial properties of Schiff base and its metal complexes were investigated against a varied range of Gram-positive and Gram-negative bacterial strains. Also, the antifungal and antioxidant assays were studied. The complex-1 shown better biological activities for antibacterial and antifungal, whereas complex-2 shown good DPPH scavenging activity.

## ACKNOWLEDGEMENT

Authors are thankful to DST-PURSE Centre, Sri Venkateshvara University, Tirupati, Andhra Pradesh, India for instrumentation facility. Authors are also thankful to Department of Biochemistry, Sri Venkateshvara University, Tirupati, A.P., India for biological activity studies.

## REFERENCES

- Prabhakaran R, Kalaivani P, Renukadevi S, Huang R, Senthilkumar K, Ramasamy K, Natarajan K, 2012. Copper Ion Mediated Selective Cleavage of C-S Bond in Ferrocenylthiosemicarbazone Forming Mixed Geometrical [(PPh<sub>3</sub>)Cu(μ-S)(2)Cu(PPh<sub>3</sub>)(2)] Having Cu<sub>2</sub>S<sub>2</sub> Core, Toward a new avenue in Copper-Sulfur Chemistry, *Inorg Chem*, 51,3525-3532.
- Prasad P, Sasmal PK, Majumdar R, Dighe R, Chakravarty AR, 2010. Photocytotoxicity and near-IR light DNA cleavage activity of oxovanadium (IV) Schiff base complexes having phenanthroline bases, *Inorg Chim Acta*, 363, 2743-2751.
- Ramachandran E, Senthil Raja D, Rath NP, Natarajan K, 2013. Role of substitution at Terminal Nitrogen of 2-Oxo-1, 2-dihydroquinoline-3-Carbaldehyde Thiosemicarbazones on the Coordination Behavior and Structure and Biological Properties of their Palladium complexes, *Inorg Chem*, 52, 1504-1514.
- Etaiw SE, Abd El-Aziz DM, Abd El-ZaherEH, AliEA, 2011. Synthesis, spectral, antimicrobial and antitumor assessment of Schiff base derived from 2-aminobenzothiazole and its transition metal complexes. *Spectrochim, Acta Part A Mol Biomol, Spectrosc*, 79, 1331-1337.
- Affan MA, Salam MA, Ahmad FB, White F, Ali HM, 2012. Organotin(IV) complexes of 2-hydroxyacetophenone-N(4)-cyclohexylthiosemicarbazone (H2dact): Synthesis, spectral characterization, crystal structure and biological studies, *Inorganica Chim Acta*, 387, 219-225.
- Bal TR, Anand B, Yogeewari P, Sriram D, 2005. Synthesis and evaluation of anti-HIV activity of isatin β-thiosemicarbazone derivatives, *Bioorg Med Chem Lett*, 15, 4451-4455.
- Panneerselvam P, Nair RR, Vijayalakshmi G, Subramanian EH, Sridhar SK, 2005. Synthesis of Schiff bases of 4-(4-aminophenyl)-morpholine as potential antimicrobial agents, *Eur J Med Chem*, 40, 225-229.
- Madappa BH, Sumathi RB, 2012. Synthesis, spectroscopic, antimicrobial and DNA cleavage studies of new Co (II), Ni (II), Cu (II), Cd (II), Zn (II) and Hg (II) complexes with naphthofuran-2-carbohydrazide Schiff base, *J Mol Struct*, 1022, 130-138.
- Jansson PJ, Sharpe PC, Bernhardt PV, Richardson DR, 2010. Novel thiosemicarbazones of the ApT and DpT series and their copper complexes, identification of pronounced redox activity and characterization of their antitumor activity, *J Med Chem*, 53(15), 5759-5769.
- Goswami TK, Gadadhar S, Roy M, Nethaji M, Karande AA, Chakravarty AR, 2012. Ferrocene-conjugated Copper(II) complexes of L-methionine and phenanthroline bases, synthesis, structure and photocytotoxic activity, *Organometallics*, 31, 3010-3021.
- Montazerzohori M, Zahedi S, Masoud Nasr-Esfahani, Naghiha A, 2014. Some new cadmium complexes, Antibacterial antifungal activity and thermal behavior, *J Ind Eng Chem*, 20 (4), 2463-2470.
- Saghatforoush LA, Aminkhani A, Ershad S, Karimnezhad G, Ghammamy S, Kabiri R, 2008. Preparation of Zinc (II) and Cadmium (II) Complexes of the Tetradentate Schiff Base Ligand2-((E)-(2-(2-(pyridine-2-yl)ethylthio)ethylimino)methyl)-4-bromophenol(PytBrsalH), *Molecules*, 13(4), 804-811.
- Tarafder MTH, Teng-Jin K, Crouse K, Ali AM, Yamin BM, Fun HK, 2002. Coordination chemistry and bioactivity of Ni<sup>2+</sup>, Cu<sup>2+</sup>, Cd<sup>2+</sup> and Zn<sup>2+</sup> complexes containing bidentate Schiff bases derived from S -benzylthiocarbamate and the X-ray

- crystal structure of bis[ S -benzyl-b- N -(5-methyl-2-furylmethylene)dithiocarbazato]cadmium(II), Polyhedron 21, 2547-44.
14. Keypour H, Dehghani-Firouzabadi AA, Rezaeivala M, Goudarziafshar H, 2010. Synthesis and characterization of Cd(II) macrocyclic Schiff base complex with two 2-Aminoethyl pendant arms, J Iran Chem Soc, 7, 820-824.
  15. Spinu C, Pleniceanu M, Tigae C, 2008. Biologically Active Transition Metal Chelates with a 2-Thiophenecarboxaldehyde-Derived Schiff Base, Synthesis, Characterization, and Antibacterial Properties, Turk J Chem, 32, 487-93.
  16. Reiss A, Florea S, Caproiu T, Stanica N, 2009. Synthesis, characterization, and antibacterial activity of some transition metals with the Schiff base N-(2-furanylmethylene)-3 aminodibenzofuran, Turk J Chem 33, 775-783.
  17. Alomar K, Landreau A, Kempf M, Khan MA, Allain M, Bouet G, 2010. Synthesis, crystal structure, characterization of zinc(II), cadmium(II) complexes with 3-thiophene aldehyde thiosemicarbazone (3TTSCH), Biological activities of 3TTSCH and its complexes, J Inorg Biochem, 104, 397-404.
  18. Omar MM, Abd El-Halim HF, Khalil EAM, 2017. Synthesis, characterization, and biological and anticancer studies of mixed ligand complexes with Schiff base and 2,2'-bipyridine, Appl Organometal Chem, 31, 01-11.
  19. Vanitha S, Prasuna KM, Saritha N, Varada Reddy A, 2019. Synthesis and Spectral Characterization of Cu (II) Complexes with 2-Hydroxy-2-Methyl Propiophenone Picolyamine and Diimine Co-Ligands, Int J Research and Analytical Reviews, 6(2),7-14.
  20. Hazra M, Dolai T, Giri S, Patra A, Dey SK, 2017. Synthesis of biologically active cadmium (II) complex with tridentate N2O donor Schiff base: DFT study, binding mechanism of serum albumins (bovine, human) and fluorescent nanowires, J Saudi Chem Soc, 21, 445-456.
  21. Mahmoud WH, Mahmoud NF, Mohamed GG, El-Sonbati AZ, El-Bindary AA, 2015. Ternary metal complexes of guaifenesin drug, Synthesis, spectroscopic characterization and in-vitro anticancer activity of the metal complexes, Spectrochim Acta A, 150, 451-60.
  22. Philip V, Sunil V, Kurup MRP, Nethaji M, 2006. Copper(II) complexes derived from di-2-pyridyl ketone N(4),N(4)-(butane-1,4-diy)thiosemicarbazone, Crystal structure and spectral studies, Polyhedron, 25(9), 1931-1938.
  23. Ibrahim MM, Abd El-Motaleb MR, Shaban Y, Mersal GAM, El-Shazly SA, Al-Juaid SS, 2017. Syntheses characterization and antioxidant activity studies of mixed-ligand copper(II) complexes of 2,2'-bipyridine and glycine, The X-ray crystal structure of [Cu(BPy)(Gly)]ClO4, J Mol Struct, 1134, 319-329.
  24. Asghar F, Badshah A, Butler I, Tabassu S, Lal BT, Tahir MN, 2016. Synthesis structural characterization in vitro bioactivities interaction with SS-DNA and DFT study of 4-chloro-3-ferrocenylaniline, Inorganica Chim Acta, 442, 46-55.
  25. Tuyen PT, Xuan TD, Khang DT, Ahmad A, Quan NV, Tu Anh TT, Anh LH, Minh TN, 2017. Phenolic Compositions and Antioxidant Properties in Bark, Flower, Inner Skin, Kernel and Leaf Extracts of Castanea crenata Sieb, et Zucc, Antioxidants, 6, 31.

#### How to cite this article

Vanitha Sanambatla, Sathish Kumar Nara, Balaji Hari, Varada Reddy Ammireddy, Saritha Nimmakayala, 2022. Antimicrobial and antioxidant studies of Schiff base, 2-(2-hydroxy-2-methyl-1-phenylpropylidene) hydrazine carbothioamide and its mixed ligand cd (II) complexes. J. Med. P<sup>r</sup>actical Allied Sci. V 11 - I 1, Pages - 4470 - 4476. doi: 10.55522/jmpas.V11I1.2451.