



## Review article

## Preliminary phytochemical and physicochemical Investigation and thin layer chromatography of Butea Monosperma flower extract

Prakash Pandey, Ashish Garg\*, Ajay Shukla

Department of Pharmaceutical Science, Guru Ramdas Khalsa Institute of Science & Technology Pharmacy, Jabalpur, Madhya Pradesh, India

**Corresponding author:** Ashish Garg, ✉ [ashish.garg071010@gmail.com](mailto:ashish.garg071010@gmail.com),

Department of Pharmaceutical Science, Guru Ramdas Khalsa Institute of Science & Technology Pharmacy, Jabalpur, Madhya Pradesh, India

© The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by-nc/4.0/>). See <https://jmpas.com/reprints-and-permissions> for full terms and conditions.

**Received** – 20 October 2014, **Revised** – 20 November 2014, **Accepted** – 25 January 2015 (DD-MM-YYYY)

**Refer This Article**

Prakash Pandey, Ashish Garg, Ajay Shukla, 2015. Preliminary phytochemical and physicochemical Investigation and thin layer chromatography of Butea Monosperma flower extract. Journal of medical pharmaceutical and allied sciences, V 4 - I 1, Pages -255 – 258. Doi: <https://doi.org/10.55522/jmpas.V4I1.0067>.

### ABSTRACT

Almost all parts of the plant namely root, leaves, fruit, stem bark, flowers, gum, young branches are used as medicine. The fresh juice of Butea monosperma is applied to ulcers and for congested and septic sore throats (1). The gum is a powerful astringent given internally for diarrhea and dysentery. The bark is reported to possess astringent bitter, pungent, alliterative, aphrodisiac and anthelmintic properties also useful in tumors, bleeding piles and ulcers. The decoction is useful in cold, cough, fever and menstrual disorders. Roots are useful in elephantiasis and in curing night blindness and other eyesight defects. Also cause temporary sterility in women. They are also used to cure boils, pimples and tumors hemorrhoids and piles. Flowers are reported to possess astringent, diuretic, depurative, aphrodisiac and tonic properties. They are used to reduce swellings. Also effective in leprosy, leucorrhea and gout. In present study Preliminary phytochemical and physicochemical Investigation and thin layer chromatography of Butea Monosperma flower extract. Various phytochemical studies are performed through which we get the knowledge about different phytoconstituent which are present in ethanolic extract of flower of Butea monosperma.

**Keywords:** Butea monosperma, Etanolic extract, Aphrodisiac, Anthelmintic, Bleeding piles, Phytochemical, Phytoconstituent.

### INTRODUCTION

The use of medicinal plants as a source for relief from illness can be traced back over five millennia to written documents of the early civilization in China, India and the Near east, but it is doubtless an art as old as mankind. The use of herbs and medicinal plant as the first medicines is a universal phenomenon. Every culture on the earth, through written or oral tradition, has relied on the vast variety of natural chemistries found in plants for their therapeutic properties. All drugs from the plant are substances with a particular therapeutic action extracted from plants. Butea monosperma (Lam.) is a sacred tree, referred to as a treasurer of the gods, and used in sacrifice related rituals. From its wood, sacred utensils are made. The flowers are offered as in place of blood in sacrifice rituals to goddess Kali. The dry stem pieces are used to make sacred fire. It is an anthropogenic tree of several castes. 'Chakradatta' mentions the use of its gum in external astringent application. The leaves

are believed to have astringent, depurate, diuretic and aphrodisiac properties. It promotes diuresis and menstrual flow. The seed is anthelmintic. The bark is also used in snakebite. When seeds are pounded with lemon juice and applied to the skin the act as a rubefacient. Arab horse dealers put one seed into each feed of corn to keep their horses in condition. The plant is used in ayurvedic, unani and siddha medicine for various ailments. Almost all parts of the plant namely root, leaves, fruit, stem bark, flowers, gum, young branches are used as medicine, food, fiber and for other miscellaneous purpose such as fish poison, dye etc. Almost 45 medicinal uses are associated with the plant and out of these claims almost half the number of claims have been scientifically studied and reported. The fresh juice of Butea monosperma is applied to ulcers and for congested and septic sore throats. The gum is a powerful astringent given internally for diarrhea and dysentery. The bark is reported to possess astringent bitter, pungent, alliterative, aphrodisiac and

anthelmintic properties. Useful in tumors, bleeding piles and ulcers. The decoction is useful in cold, cough, fever and menstrual disorders. Roots are useful in elephantiasis and in curing night blindness and other eyesight defects. Also cause temporary sterility in women. Also applied in sprue, piles, ulcers, tumors. Leaves have astringent, tonic, diuretic and aphrodisiac properties. They are also used to cure boils, pimples and tumors hemorrhoids and piles. Flowers are reported to possess astringent, diuretic, depurative, aphrodisiac and tonic properties. They are used to reduce swellings. Also effective in leprosy, leucorrhea and gout. In 1938, TLC was developed by Ismailoff and Schraiber. An adsorbent is coated on a glass plate which serves as a stationary support on which the mobile phase percolates and develops the chromatogram. This method is simple, rapid in separation and sensitive. The speed of separation is fast and it is easy to recover the separated compounds from plate.

## MATERIALS AND METHOD

### Materials

Fresh flowers about 150gm of *Butea monosperma* were collected in the flowering season is from March-April. Ethanol (95%) was purchased from CDH chemical laboratory, New Delhi, India, Silica Gel "G", Toluene, Ethyl Acetate and Vanillin-Sulphuric Acid Reagent were purchased from Spectrochem Mumbai, India. All other chemicals used were of analytical reagent grade and were utilized as they received.

### Method

#### Preparation of Extract

The extraction process of flower of *Butea monosperma* was done by maceration process

In this process initially 150gm of dried flowers were weighed and put in 250ml conical flask then 200ml of extraction fluid (ethanol) was added on this and the fluid was poured on dried flowers of *Butea monosperma* previously weighed. The mixture was then kept for three days in a tightly sealed flask at room temperature protected from sunlight. The mixture was shaken several times daily. The mixture was filtered and filtrate was collected. The extract was concentrated under atmospheric pressure. The extract prepared was then transferred to suitable container and the further experiment work was started.

#### Standardization Parameters

#### All the parameters performed as per W.H.O. (WORLD HEALTH ORGANIZATION)

#### Guidelines.

#### Determination of ash value

A 3gm of the grounded air dried material was taken in previously ignited and tarred crucible. The material was then spread in an even layer and ignited in by gradually increasing the heat to 500-600 until its white indicating an absence of carbon. Then it was cool in desiccators for 30min and then weighed. Then content of total ash was calculated in mg/gm of dried material. Acid Insoluble Ash.

To the crucible containing the total ash 25ml of hydrochloric acid was added and cover with a watch glass and then boiled gently for 5min. Watch glass was then rinsed with hot water and this liquid was

added to crucible. The insoluble material was collected on an ash less filter paper and then washed with hot water until the filtrate was neutral. The filter paper containing the insoluble matter was transferred to original crucible dry on a hot plate and was ignited to a constant weight. The residue was allowed to cool in suitable desiccators for 30min and then weighed. The content of acid insoluble ash was collected in mg/gm of dried material [2].

#### Water Soluble Ash

To the crucible containing the total ash 25ml of water added and then boiled for 5min. The insoluble matter was collected on an ash less filter paper. Then it was washed with hot water and ignited it in a crucible for 15min at a temperature not exceeding 450. The weight of this residue in mg is subtracted from the weight of total ash then the content of water soluble ash was calculated in mg/gm of air dried material.

#### Determination of Extractable Matter (Extractive Value)

4gm of coarsely powdered air dried material accurately weighed in a glass stoppered conical flask. It was then macerated with 100ml of ethanol for 6 hours. Then it was allowed to stand for 18 hours. It was filtered and 25ml of the filtrate was transferred to a tared bottom dish and evaporated to dryness on a water bath. It was dried at 105 for 6 hours. It was cooled in desiccators for 30min and then it was weighed. Content of extractable matter was calculated in mg/gm of air dried material.

#### Determination of Loss on Drying

Determination of loss on drying is also known as gravimetric determination. 3gm of material was accurately weighed place in a previously dried and tarred weighing bottle. The sample was dried by both the methods. In an oven at 100-105°C and in a desiccators over phosphorous pentoxide under atmospheric pressure or reduced pressure and at room temperature. It was dried till 2 consecutive readings don't differ by 5 mg. Then at last loss of weight in mg/gm of air dried material was calculated [3].

#### Determination of Swelling Index

Simultaneously three determinations for the given material. The material was previously reduced to required fineness. 3gm of the material was accurately weighed. It was placed into a 25ml glass stoppered measuring cylinder. 25ml of water was added and the mixture was shaken thoroughly for

1hour. It was allowed to stand for 3 hours at room temperature. The volume occupied by the material was measured. The mean value of the individual determinations was carried out [4].

#### Preliminary Phytochemical Screening Test for Alkaloids

Dragendorff's test: - In 2-3ml of filtrate, few drops of previously prepared Dragendorff reagent was added.

Mayer's test: - In 2-3ml of filtrate, few drops of Mayer reagent was added.

Hager's test: - In 2-3ml of filtrate, few drops of Hager reagent

was added.

Wagner's test: - In 2-3ml of filtrate, few drops of wagner reagent was added.

#### Test for Flavonoids

In 2-3ml of filtrate, Lead Acetate solution was added.

In Few drops of filtrate, Sodium Hydroxide solution was added

Shinoda Test: - In Few drops of filtrate, 5ml of Ethanol, Few drops of Conc. HCl and Magnesium Turnings were added.

#### Test for Tannins

In 2-3ml of alcoholic extract, Ferric Chloride solution was added.

In 2-3ml of alcoholic extract, Lead Acetate solution was added.

In 2-3ml of extract, Acetic Acid was added.

In 2-3ml of extract, dilute Nitric Acid was added.

In 2-3ml of ethanolic extract, Potassium Permanganate solution was added.

#### Test for Steroids

Salkowski Test: - In 2ml of extract, 2ml of Chloroform, 2ml of Conc. Sulphuric Acid were added and shaken well.

Lieberman Burchard Test: - In 2ml of extract, 2ml of Chloroform, 2ml of Conc. Sulphuric Acid and 2ml of Acetic Anhydride Solution were added.

Lieberman Test: - In 3ml of extract, 3ml of Acetic Anhydride and few drops of Conc. Sulphuric Acid were added.

#### Test for Carbohydrate

Test for Reducing Sugar

Fehling Test: - In Few drops of extract, 1ml of Fehling solution A and B was added

#### Test for Pentose

Aniline Acetate Test: - The test solution (flower extract) was boiled and treated with filter paper soaked in Aniline acetate solution was added.

#### Test for Hexose

Cobalt Chloride Test: - In 3ml of test solution, 2ml of Cobalt Chloride solution was added and then boiled and cooled after that few drops of Sodium Hydroxide solution was added.

#### Test for Glycoside

#### Test for Saponin

Foam Test: - Drug extract was shaken vigorously with water.

Test for anthraquinone Glycoside: - In 3ml of extract, dilute sulphuric acid and equal volume of benzene was added and shaken [5].

#### Thin Layer Chromatography

##### Adsorbent - Silica Gel "G"

Mobile Phase - Toluene : Ethyl Acetate (9:1) Detecting Agent - Vanillin-Sulphuric Acid Reagent Sample For Detection- Ethanolic Extract of *Butea monosperma* Flower TLC plates were prepared by mixing the adsorbent (silica gel "G") with a small amount of inert binder like calcium sulfate (gypsum) and water. This mixture was spread as thick slurry on glass slide. The TLC plates were dried and

activated by heating in an oven for 30min at 110 °C. The thickness of the adsorbent layer is typically around 0.1–0.25mm for analytical purposes. A small spot of solution containing the sample (Extract) is applied to a TLC plate, about 1cm from the bottom edge. The solvent is allowed to completely evaporate off. A small amount (10ml) of Mobile Phase (Toluene:Ethyl Acetate) was poured in to a glass beaker and TLC plate was put into the chamber, so that its bottom touches the solvent, and the plate lies on the chamber wall and reaches almost to the top of the container. The container was closed with a cover glass and plate was left for a few minutes to move the mobile phase on TLC plate by capillary action. When the mobile phase was reached on the top of the TLC plate and separation was take place in the separating chamber, the plate was removed and dried. After dryness of the TLC plate the detecting agent (Vanillin-Sulphuric Acid Reagent) was sprayed and observed the plate under UV lamp. Then  $R_f$  Value was calculated [6].

The Standardization Parameters investigation of ethanolic extract of flower of *Butea monosperma* can be performed under the following methods like Total Ash Value, Extractive value, Swelling index and under

loss on drying categories. The data was obtained of following estimation was satisfactory and shows that the phytochemical agents are present in the flower extract in effective quantity as well as in quality. The results of Standardization Parameters.

The phytochemical investigation ethanolic extract of flower of *Butea monosperma* can be performed under the following tests like Alkaloids, Flavonoids, Tannins, Steroid, Carbohydrates, and Glycoside categories. The result was obtained of following estimation was satisfactory and shows that the Alkaloids, Flavonoids, Carbohydrate, Tannins and Steroids like chemical constituents are present in the flower extract in effective quantity and quality. The results of phytochemical study.

Thin layer chromatography was carried out using Toluene: Ethyl Acetate (9:1). After running the solvent,  $R_f$  Values were calculated for the obtained two spot. On the basis of  $R_f$  Values (0.31 and 0.65), the compounds were identified as Butrin and Isobutrin [7, 8].

#### CONCLUSION

The flower of *Butea monosperma* has many uses which include Anthelmintic, dysentery, cold, cough, fever, menstrual disorders, tumors, bleeding piles and ulcers, night blindness and other eyesight defects. In present study Preliminary phytochemical and physicochemical Investigation and thin layer chromatography of *Butea Monosperma* flower extract. Various phytochemical studies are performed through which we get the knowledge about different phytoconstituent which are present in ethanolic extract of flower of

*Butea monosperma*.

## REFERENCES

1. Igwe SA, Akunyili DN, 2005. Analgesic Effects of Aqueous extracts of the leaves of *B. Pinnatum*. *Pharmaceutical Biology*. 43(8), Pages 658- 661.
2. Nooman A, Khalaf Ashok K, Shakya K, 2008. Antioxidant Activity of Some Common Plants. *Turk J Biol*. 32, Pages 51-55.
3. Beever D E Rumen, 1993. Function .in quantitative aspects of ruminant digestion and metabolism, forbes J.(Eds), Wallingford CAB International. Pages 187-215.
4. Callaway TR, Martin SA, Wampler JL, Hill GM, 1999. Malate content of forage varieties commonly fed to cattle. *Journal of dairy science*. 80, Pages 1651-1665.
5. Rao NK, Pundir RPS, Van Der Maesen LJG, 1980. Inheritance of some qualitative characters in chickpea (*Cicer arietinum* L.). *Indian Acad. Sci. (Plant Sci.)*. 89(6), Pages 497-503.
6. Shah Isha, Bhatt Shailendra, Yadav Alpesh, 2014. Enhancement of solubility and dissolution of nebivolol by solid dispersion technique. *Int J of Pharmacy and Pharmaceutical Sciences*. 6(7), 566-571.
7. Ojewole JAO. 2005. Anti- nociceptive, Anti-inflammatory and Antidiabetic effect of *Bryophyllum pinnatum* (Crassulaceae) leaf aqueous extract. *J of Ethano pharmacology*. 99, Pages 13–9
8. Gaind KN, Gupta RL, 1974. Identification of waxes from the leaves of *Kalanchoe pinnata*. *Planta medica*, 23, Pages 193–7.