



Journal of Medical Pharmaceutical and Allied Sciences

Journal homepage: www.jmpas.com CODEN: JMPACO

Research article

Standardization of Marketed Herbal Product Kawach Beej Churna

Rajmammad Husen Shaikh*, Mojsin J Jamadar

Department of Pharmaceutics Appasaheb Birnale College of pharmacy, Sangli, Maharashtra, India

Corresponding author: Rajmammad Husen Shaikh, [✉ rajshaikh71@gmail.com](mailto:rajshaikh71@gmail.com),

Department of Pharmaceutics Appasaheb Birnale College of pharmacy, Sangli, Maharashtra, India

© The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by-nc/4.0/>). See <https://jmpas.com/reprints-and-permissions> for full terms and conditions.

Received – 10 October 2015, Revised – 20 November 2015, Accepted – 25 February 2016 (DD-MM-YYYY)

[Refer This Article](#)

Rajmammad Husen Shaikh, Mojsin J. Jamadar, 2016. Standardization of Marketed Herbal Product Kawach Beej Churna. Journal of medical pharmaceutical and allied sciences, V 5 - I 1, Pages -293 – 299. Doi: <https://doi.org/10.55522/jmpas.V5I1.0077>.

ABSTRACT

Herbal medicines are the most ancient form of health care known to human beings. It consists of plant or its part to treat injuries, disease or illnesses and is used to prevent and treat diseases and ailments or to promote health and healing. It is a drug or preparation made from a plant or plants and used for any of such purposes. Now a day, approximately 70% of "synthetic" medicines are derived from plants. Organoleptic evaluation revealed that Kawach churna MK-3 was observed to be whitish brown in color. Results of Extractive value demonstrated that water soluble extractive value and alcohol soluble extractive value of product MK-3 was lesser as compared to products MK-1, MK-2 and LB-1. Phytochemical studies exhibited presence of Alkaloids, Steroids, Saponins, and Tannins in all the four products. Ash value determination revealed that the acid insoluble ash of all the products was within range < 1%, but acid insoluble ash of product MK-3 was more as compared to products MK-1, MK-2 and LB-1, which indicates presence of sand or earth material in product MK-3. Moisture content of all the products were within acceptable ranges < 5 % w/w. But as product MK-3 contained more moisture it is more prone to microbial contamination and enzymatic degradation. When all the four products were subjected for Microbial Limit test, all the compounds demonstrated growth of non-pathogenic micro-organisms but growth of the pathogenic micro-organisms was not observed. Product MK-2 may be better than products MK-1 and MK-3 in view of safety and efficacy, as it contains higher quantity of active markers. TLC showed yellow spot having Rf value 0.73 in n-butanol: Glacial Acetic Acid: Water (8:2:2) (v/v/v) solvent system. UV and IR spectra of isolated and standard marker compound (Levodopa) showed similar results.

Keywords: Asthma control therapy, Drug utilization study, Pharmacist observation.

INTRODUCTION

Medicinal plants are the nature's gift to human being to make disease free healthy life. It plays a vital role to preserve our health. Plants and their products are being used as a source of medicine since long. They provide basic compounds affording less toxic and more effective drug molecules. Popularity among the common people increased the usage of medicinal plants/herbal products. World Health Organization (WHO) has defined herbal medicines as finished, labeled medicinal products that contain active ingredients, aerial or underground parts of the plant or other plant material or combinations. World Health Organization has set specific guidelines for the assessment of the safety, efficacy, and quality of herbal medicines. In drug industry, herbs are used for following reasons. Source of phyto-pharmaceuticals: To isolate pure compounds. Herbal raw materials: To formulate herbal, Ayurvedic, Homeopathic and

Unani medicines. Medicinal herbs: To prepare plant extracts and single ingredient powders. Dietary supplements: Sources of vitamins and enzymes. The emergent use of herbal drug by the human is forcing moves to evaluate the health claims of these agents and to develop standards of quality, purity, safety and efficacy of the drug. The mostly herbal drugs are effective but lack of standardization. So there is need to develop standardization parameters.

Herbal medicines make up an important part of the trend toward alternative medicine.

In many countries herbal drugs are lunched into the market without standard scientific evaluation.

Herbal Drugs

Traditionally, herbal drugs have played a significant role in the management of both minor and major medical illnesses. Herbal drugs

means a "dosage form consisting of one or more herbs or processed herb in a specified quantities to provide specific nutritional, benefits, and/or other benefit meant for use to diagnosis treat, mitigate diseases of human beings or animal and/or alter the structure or physiology of human beings or animals". Herbal drugs are obtained by whole plant, cut plant part, powdered plant part, extract, essential oil, expressed juices and plant parts to treatment such as distillation, extraction, expression, fractionation, purification and concentration. Herbal medicinal product is any medicinal product, exclusively containing as active substances one or more herbal substances or one or more herbal preparations, or one or more such herbal substances in combination with one or more such herbal preparations. Herbal substances are mainly whole, fragmented or cut plants, plant parts, algae, fungi and lichen in an unprocessed, usually dried form but sometimes fresh. Certain exudates that have not been subjected to a specific treatment are also considered to be herbal.

Advantages of Herbal Drugs

They have low risk of side effects.

They have more effective than any synthetic drug.

They have lower cost.

They have large amount of availability.

They have better patient tolerance as well as acceptance.

They have large therapeutic activity.

The medicinal plants have renewable source of cheaper medicines.

Many natural medicines are being shown to produce better results than drugs or surgery without the side effects.

They have a strong traditional or conceptual base and the potential to be useful as drugs in terms of safety and effectiveness whereas modern medicines have a very strong experimental basis for their use but are potentially toxic.

They can deliver mixtures of multi- functional molecules with potentiating and synergistic effects.

They are well suited for long-term disease prevention in an era of genetic testing and increased life expectancy.

Herbal products have better compatibility with the human body.

A prolong therapy dose not showed any side effect by use of most of the herbal products.

Herbal products have long history of use and better patient tolerance as well as acceptance.

The cultivation and processing of medicinal herbs and herbal products is environmental friendly.

Medicines are available in both crude as well as a pure chemical form.

They also provide additional vehicles for delivering health and wellness.

Standardization 1.3.1General Introduction

Traditional medicine implies knowledge and practice of herbal therapy for the prevention, diagnosis and elimination of physical, mental, or social imbalance. The costs for health care are rising at an alarming rate throughout the world. At the same time, the world market for

phytopharmaceuticals is growing steadily. Standardization involves adjusting the herbal drug preparation to a defined content of a constituent or a group of substances with known therapeutic activity by adding excipients or by mixing herbal drugs or herbal drug preparations. Standardization of herbal drug is necessary to confirm its identity and to determine its quality, purity, safety, content, physical, chemical, biological properties. But the most important challenges faced by these herbal drugs are the lack of complete standardization by physiochemical. The standardization of herbal drugs,

Subfamily: Faboideae. **Tribe:** Phaseoleae **Genus:** Mucuna.

Species: M. pruriens.

Botanical Name: Mucuna pruriens

Other Names: Mucuna pruriens, Velvet Bean, Cowhage, Cow-itch, and Buffalo bean, Kaunch, Atmagupta, Kapikachhu. Mucuna pruriens or Kaunch is one of the popular medicinal plant of India. It is widespread over most of the subcontinent and is found in bushes and hedges and dry-deciduous, low forests throughout the plains of India. Parts of Mucuna possess valuable medicinal properties and there is a heavy demand of Mucuna in markets. The main constituent of Mucuna pruriens seeds is L-dopa or Levadopa which acts as a natural dopamine. The seeds of Mucuna pruriens are high in protein (20- 29 %), lipids (6-7%), dietary fiber (8-10%), ash (3%), carbohydrates (50-60%) and minerals. Also they are extremely rich in alkaloids, saponins, and sterols [1].

Main use of the plant are following:

It contain L-Dopa which is used as anti-Parkinson's, help increases testosterone

Aphrodisiac - increases libido and it is very good for impotency and erectile dysfunction

It is used to stimulate growth of hormone - anabolic/androgenic •It help reduces spasms and relieves pain - analgesic/ antispasmodic

Help lowers blood sugar and blood pressure (hypo tensive)

It increases urination (uterine stimulant), help reduces fever (febrifuge) and it is a good cough suppressant

It help lowers cholesterol (hypocholesterolemic)

It act as a weight loss aid

It reduces inflammation - anti- inflammatory

Mucuna is an annual twining plant. Leaves are trifoliate, gray-silky beneath; petioles are long and silky, 6.3-11.3 cm. Leaflets are membranous, terminal leaflets are smaller, lateral very unequal sided. Dark purple flowers (6 to 30) occur in drooping racemes.

Fruits are curved, 4-6 seeded. The longitudinally ribbed pod, is densely covered with persistent pale brown or grey trichomes that cause irritating blisters. Seeds are black ovoid and 12 mm long. Mucuna seeds (as well as the seeds of all Mucuna species) contain high concentration of L-dopa (7-10%) - a direct precursor of the neurotransmitter dopamine. The hairs of the seed pods contain the phytochemical mucunain, the very constituent that irritates the skin when in contact. The serotonin is also has been found

in the pod (as well as in the leaf and fruit).

Other important Mucuna properties

It helps prevent or eliminate kidney stones (antilithic)

It helps kill parasites (antiparasitic) and expels worms (vermifuge)

It expels gas (carminative)

It is a menstrual stimulant

It is a central nervous system stimulant

Benefits of Mucuna pruriens in nervous system disorder

It is used as a nervine tonic for nervous system disorders.

Because of the high concentration of L-dopa in the seeds. It has been studied for its possible use in Parkinson's disease. It was found to slow the progression of Parkinson's symptoms (such as tremors, rigidity, slurring, drooling, and imbalance), and to have none of the side-effects of the current pharmaceutical L-dopa.

Objective

Evaluation of procured product as per the various standardization parameters, which includes.

To carry out organoleptic parameters.

To carry out physico-chemical parameters.

To determine solvent extractive values: includes water, alcohol soluble extractives.

To determine ash values: includes total ash, acid insoluble ash and water soluble ash.

To determine the moisture content.

To carry out microbial limit tests.

To carry out phytochemical tests.

Extraction and isolation of the marker compound from marketed herbal medicine.

To compare isolated marker compound with pure substances (marker compounds) using chromatographic and spectroscopic methods.

Experimental Work

Organoleptic evaluation

Colour, Odour and Taste - All the four samples were analyzed for their morphological characteristics. All the samples were visually analyzed; a small portion of the drug was taken, slowly and repeatedly inhaled the air over the material and examined the odour. A small portion of drug was taken on the tongue and the taste of formulation was found.

Determination of solvent extractive values

Solvent extractive value is the amount of active constituent in a specified weight of medicinal plant material when extracted with specific solvent. The extraction of any crude drug with a particular solvent yields a solution containing different phytoconstituents. The composition of these phytoconstituents in that particular solvent depends upon the nature of the drug and solvent used. The solvent extractive value was determined by following methods to measure the water soluble extractive value and alcohol soluble extractive value using water and ethanol as solvent for extraction respectively.

Determination of water soluble extractives

5 g of churba of each manufacturer were weighed and taken

into the four conical flasks in which 100 ml of water were added separately and allowed to macerate for 24 h, shaking frequently for first 6 h then allowed standing for 18 h. Which was then filtered rapidly and the 25 ml of the filtrate was allowed to evaporate to dryness in a tared 250 ml beaker at 105°C, the difference in weight of the beaker is indication of water soluble extractive value of that drug with respect to the amount of drug (5 g) taken for extraction.

Determination of alcohol soluble extractive value:

5 g of churba of each manufacturer were weighed and taken into the conical flask in which 100 ml of ethanol were added separately and allowed to macerate for 24 hours, shaking frequently for first 6 hours, then allowing to stand for next 18 hours, which was then filtered rapidly and 25 ml of the filtrate was allowed to evaporate to dryness in a tared 250 ml beaker at 105°C. The difference in weight of the beaker is an indication of alcohol soluble extractive value of that drug with respect to the amount of drug (5 g) taken for extraction.

Determination of ash values

The ash of any organic material is composed of their non-volatile inorganic components. Controlled incineration of crude drug results in an ash residue consisting of an inorganic material (metallic salts and silica) in certain drugs, the percent variation of the weight of ash from sample to sample is very small and any marked difference indicates a change in quality. The ash value can be determined by three different methods to measure the total ash, the acid insoluble ash and the water soluble ash [2].

Determination of total ash content

The total ash was determined by placing 2 g of accurately powdered material into the tared crucible. Then samples were placed in muffle furnace and the samples were ignited by gradually increasing the heat up to 500- 600°C until it is white, indicating the absence of carbon. The crucible was kept in desiccator and weighed. Initially carbon free ash was not obtained thus the cooled crucible was removed and the residue was moistened with about 2 ml of water, dried on water bath, and ignited to constant weight. The residue was allowed to cool in desiccator for 30 minutes. Content of total ash was calculated with reference to the amount of sample taken.

Determination of acid insoluble ash value

The total ash which was obtained in the previous step was boiled with 25 ml of 2 M HCl for 5 minutes, then it was filtered through ash-less filter paper, the insoluble matter was collected in a previously tared crucible, then the residue was washed with hot water, then ignited in muffle furnace at about 450°C for 30 minutes, cooled in desiccator and weighed. The percent of acid insoluble ash was calculated with reference to the amount of sample taken.

Determination of water-soluble ash

In the crucible containing total ash, 25 ml of water was added and boiled for 5 minutes. The insoluble matter was collected on an ash less filter paper. This was then washed with hot water and ignited in a

crucible for 15 minutes at a temperature not exceeding 450°C separately. Then the weight of this residue was subtracted from the weight of total ash. Finally the content of water soluble ash with reference to the amount of sample taken was calculated.

Determination of moisture content

Moisture is an inevitable component of crude drug, which must be eliminated as far as practicable. Drying of crude drug is important during collection of drug and is also important for preservation, preventing hydrolytic degradation of active constituents and for easy size reduction of crude drug. Excess moisture or insufficient drying is responsible for spoilage of drug due to growth of microbes. Therefore drying process should reduce the moisture content of drug below the critical level. Moisture content determination is used to check the total water content in given weight of crude drug. Excess moisture content in a crude drug sample suggests not only that the purchaser could be paying a high price due to unwanted water, but also that the drug has been incorrectly prepared or subsequently incorrectly stored. Method of determination of moisture content includes the loss on drying. It can be carried out either by heating at 100°- 105°C or in a desiccator over phosphorus pentoxide under atmospheric pressure at room temperature for specific period of time.

Method: - Loss on drying, Azeotropic distillation method and

Karl fisher

method use for moisture content determination in crude drug.

Loss on drying

Loss on drying is the loss of mass expressed as percent w/w.

To estimate the loss on drying 2-5 g of air dried drug is accurately weighed in dried and tared flat weighing bottle. The substance is dried to a constant mass or for the prescribed time as specified. 5 g of sample was accurately weighed in separate dried and tared petridish. Sample was placed in an oven, and drying of sample was carried out at 105°C until a constant mass of the sample was not observed. The difference in weight before and after heating in oven was recorded and loss on drying was expressed as percent w/w.

Determination of physical parameters

Determinations of angle of repose and flow rate

Good flow properties are critical for the development of any pharmaceutical product including powder formulations. It is essential that an accurate assessment of flow properties to be made as early in the development process so that an optimum formulation can be quickly identified. Interparticle forces between particles as well as flow characteristics of powders are evaluated by angle of repose.

Angle of repose is defined as the maximum angle possible between the surface of pile of sample and horizontal plane. Angle of repose is affected by presence of moisture, size and shape of sample, percent fines, and amount of glidant and lubricant. More the moisture in sample, more it becomes sticky. Similarly smaller the size more is the flow of powder. Angle of repose is a helpful parameter to monitor quality

of powdered or granular pharmaceutical formulations. For good flowing powders angle of repose should be less than 30°.

Method

Take a clean and dry funnel with a round stem of 20 to 30 mm diameter with flat tip and attached it to the burette stand.

Place a graph paper sheet below the funnel, on clean and dry platform.

Adjust the distance between lower tip of the funnel and sheet to some specified height.

Gently pour sample in funnel from top till a heap of powder forms and touches of lower tip of the funnel.

Repeat the procedure four times to obtain average reading.

Find out average diameter and radius of the each drawn circle.

Angle of repose Calculation

Radius determination - $r = \text{diameter of heap (d)} / \text{height of heap (h)}$

Average radius determination - $r = r_1 + r_2 + r_3 + r_4 / 4$

Angle of repose (θ) - $\tan^{-1} (h/r)$

Flow rate calculation = mass of powder / time required for flow = W_2/t (gm/sec)

Determination of Bulk and Tap Density and Compressibility Index

Requirement: - Graduated measuring cylinder and bulk density apparatus and weighing balance.

Method

Weigh accurately 25 g of powder (W1).

Place it in dried graduated measuring cylinder and note the volume as V1 ml.

Place the cylinder containing sample in bulk density apparatus for 100 tapping and operate it. Record the volume occupied by the powder as V2 ml.

Bulk and tap density Calculation

Bulk density = Mass / Bulk volume = W_1 / V_1 (g/ml)

Tap density = Mass / Tap volume = W_1 / V_2 (g/ml)

Compressibility index = tap density - Bulk density / tap density

Microbial limit tests

Preliminary testing

The methods given are invalid unless it is demonstrated that the test specimens to which they are applied do not, of themselves, inhibit the multiplication under the test condition of microorganisms that can be present.

Preparation of sample for preliminary testing

1gm of sample was taken to prepare 50ml of fluid lactose medium. But this medium was not suitable for sufficient growth of *Escherichia coli* and *Pseudomonas aeruginosa*. Hence the relevant medium was prepared by taking 1gm of sample to prepare 100ml of fluid lactose medium. Prior to doing the test, diluted specimen of substance being examined was inoculated with separate viable cultures of *Escherichia coli*, *Salmonella* species, *Pseudomonas aeruginosa* and *Staphylococcus aureus*. This was done by adding 1ml of 24 hours cultures of the microorganisms to the first dilution (in buffer solution pH 7.2, fluid lactose medium) of the test material.

Initially *Escherichia coli* and *Pseudomonas aeruginosa* organisms failed

to grow in the relevant medium. The procedure was modified by increasing the volume of diluents with the quantity of test material remaining the same, i.e. 1gm of sample was taken to prepare 100 ml of fluid lactose medium. After this modification growth of the strains was observed.

Tests for specified micro- organisms

Test for *E. coli*

1 g of sample was added to the MacConkey broth and incubated at 43- 450C for 24 hours. A subculture with the same media was made and incubated at the same temperature for the same time. Growth of red, generally no mucoid colonies of gram negative rods, sometime surrounded by a reddish zone of precipitation, indicates the possible presence of *E. coli*. Therefore, the material will pass the test if no such colonies are grown.

Test for *S. typhi*

The solution of the pretreated sample was prepared and incubated as above for 24 hours. 10ml of the enrichment medium was transferred to 100ml of tetrathionate bile brilliant green broth and incubated at 42-430C for 24 hours. Subculture was prepared by streaking method, with the agar media like, Xylose lysine agar. Incubated at 35-370C for 24 hours and look out for well-developed red with or without black centered colonies, indicates the possible presence of *S. typhi*. Therefore, the material will pass the test if no such colonies are grown.

Test for *P. aeruginosa*

The 1 gm of sample was pretreated with buffered sodium chloride peptone solution, pH 7.0. Inoculated 100ml of soyabean casein digest broth the solution, suspension or emulsion was obtained. After mixing it was incubated at 35-370C for 48 hours. Subculture was prepared on a Cetrimide agar plate and incubated at the same condition. If there is no growth of the microorganisms then the material passes the test. If growth of colonies of gram negative rods occurs usually with greenish fluorescence, then it indicates the presence of the *Pseudomonas* species.

Test for *S. aureus*

The pretreated material of all four samples were used and inoculated. It has to be incubated in Vogel- Johnson agar at 35-370C for 24 h. If there is no growth of the microorganism then the material passes the test. The presence of *Staphylococcus aureus* will be detected with the presence of black colonies of gram positive cocci often surrounded by a clear zone.

Phytochemical tests

The alcoholic extracts of churna were subjected to qualitative chemical investigation. The following procedures were adopted to test the presence of various chemical constituents in the extracts.

Test for alkaloids

Mayer's test: Treat the extract with Meyer's reagent (potassium mercuric iodide) gives cream colored precipitate.

Wagner's test: Treat the extract with Wagner's reagent (iodine in

potassium iodide) gives brown precipitate.

Dragendorff's test: Treat the extract with Dragendorff's reagent (potassium bismuthiodide) gives reddish brown precipitate.

Hager's test: Treat the extract with Hager's reagent (saturated solution of picric acid) gives yellow colour.

Test for steroids

Salkowski test: Treat the extract with few drops of concentrated sulphuric acid; red colour at the lower layer indicates presence of steroids.

Libermann-Burchard test: Treat the extract with few drops of acetic anhydride, boil and cool. Then added concentrated sulphuric acid from the side of the test tube, brown ring is formed at the junction of two layers and upper layer turns green which indicates presence of steroids.

Test for flavonoids

Shinoda test: To the extract, add few magnesium turnings and concentrated hydrochloric acid dropwise, crimson red or green to blue colour appears after few minutes which indicates presence of flavonoids.

Alkaline reagent test: To the extract, add few drops of sodium hydroxide solution, intense yellow colour is formed which turns to colourless on addition of few drops of dilute acid indicates presence of flavonoids.

Test for saponins

Foam test: Small quantity of churna treated with water and shaken for sometime which shows the formation of foam which remained stable for 15 minutes.

Bromine water test: The test solution treated with bromine water separately, which gives yellow precipitate.

Test for tannins

Ferric chloride test: Test solution with few drops of Ferric chloride solution gives dark colour.

Gelatin test: Test solution when treated with gelatin solution forms white precipitate.

Extraction and isolation of marker constituents from marketed Kawach churna

Isolation of Levodopa from Marketed Kawach Churna

The dried, milled seeds (100 g) were defatted with acetone (300ml) by shaking for 48 h at room temperature. Defatted material was extracted with water- ethanol (1:1) with 0.1% ascorbic acid by shaking overnight. The residue was removed by filtration and filtrates were pooled and concentrated. The concentrate after crystallization yielded crude L dopa, which on further recrystallization in hot water gave pure crystals.

For identification, isolated samples were subjected to Qualitative tests, IR spectroscopy were determined and compared with standard sample of Levodopa.

Qualitative Tests for Levodopa

Dissolve 2 mg in 2 ml of water and add 0.2 ml of Ferric chloride solution Green colour develops which changes in bluish violet on addition of 0.1 g of hexamine.

IR spectroscopy

The IR analysis of standard levodopa was carried out using

pressed pellet technique [f, and presence of probable functional groups were interpreted.

Similarly the isolated levodopa of all the four samples were analyzed for their IR spectra, and the IR spectra of the isolated levodopa of all the four samples were Compared (overlaid) with that of standard levodopa

Qualitative and Quantitative Estimation of isolated levodopa by spectroscopic analysis

For Estimation, isolated levodopa was subjected to UV spectroscopy and results were compared with standard levodopa (RS).

UV spectroscopy

Determination of λ max for standard levodopa (RS) and isolated levodopa

Preparation of stock solution of standard levodopa (RS) and their dilutions

An accurately weighed quantity of about 10 mg of standard levodopa was taken in

100 ml volumetric flask dissolved in sufficient quantity of 0.1 N HCL then sonicated for 15 min and diluted up to

100 ml of 0.1 N HCL to yield a concentration 100 μ g/ml. Further dilutions were made, 0.5 ml of stock solution was separately transferred to 10 ml volumetric flask and volume was made up to 10ml in volumetric flask with 0.1 N HCL to yield a concentration of 5 μ g/ml likewise 1 ml of stock solution in

10 ml of 0.1 N HCL, 1.5 ml of stock solution in 10 ml of 0.1 N HCL, 2 ml of stock solution in 10 ml of 0.1 N HCL and 2.5 ml of stock solution in 10 ml of

0.1 N HCL to yield a concentration 10 μ g/ml, 15 μ g/ml, 20 μ g/ml and 25 μ g/ml.

The prepared dilutions of standard levodopa were scanned separately in a wavelength range of 200-400 nm against

0.1 N HCL as blank to get maximum absorption for the standard levodopa. And above same procedure was followed for preparation of stock solution of isolated levodopa and their dilutions. The prepared dilutions of isolated levodopa were scanned separately in a wavelength range of 200-400 nm by using UV spectrophotometer against 0.1 N HCL as blank to get maximum absorption for the isolated levodopa [3,4].

RESULT AND DISCUSSION

Organoleptic Evaluation

Organoleptic properties of Kawach churna of different manufacturers were similar i.e. Brown colour, characteristic odour and sweet taste. The only difference was that Kawach churna of MK-3 was whitish brown in colour.

Extractive Value

The water soluble extractive of kawach churna of different manufacturers MK-1, MK-2, MK-3 and LB-1 was found to be 31.5%, 32.18%, 30.68% and 32.62% w/w respectively.

The alcohol soluble extractive of kawach churna of different manufacturers

MK-1, MK-2, MK-3 and LB-1 was found to be 15%, 15.38%, 14.94% and 15.7% w/w respectively. The total ash value of kawach churna of different manufacturers MK-1, MK-2, MK-3 and LB-1 was found to be 4.04%, 3.9%, 4.06% and 3.16% w/w respectively. The water soluble ash value of kawach churna of different manufacturers MK-1, MK-2, MK-3 and LB-1 was found to be 1.68%, 0.62%, 1.72% and 1.27% w/w respectively. The acid insoluble ash value of kawach churna of different manufacturers MK-1, MK-2, MK-3 and LB-1 was found to be 0.53%, 0.39%, 0.58% and 0.36% w/w respectively [5, 6].

CONCLUSION

Organoleptic evaluation revealed that Kawach churna MK-3 was observed to be lightish brown in color. Results of Extractive value demonstrated that water soluble extractive value and alcohol soluble extractive value of product MK-3 was lesser as compared to products MK-1, MK-2 and LB-1. Phytochemical studies exhibited presence of Alkaloids, Steroids, Saponins, and Tannins in all the four products. Ash value determination revealed that the acid insoluble ash of all the products was within range < 1%, but acid insoluble ash of product MK-3 was more as compared to products MK-1, MK-2 and LB-1, which indicates presence of sand or earth material in product MK-3. Moisture content of all the products were within acceptable ranges < 5 % w/w. But as product MK-3 contained more moisture it is more prone to microbial contamination and enzymatic degradation. When all the four products were subjected for Microbial Limit test, all the compounds demonstrated growth of non-pathogenic micro-organisms but growth of the pathogenic micro-organisms was not observed. Product MK-2 may be better than products MK-1 and MK-3 in view of safety and efficacy, as it contains higher quantity of active markers. TLC showed yellow spot having Rf value 0.73 in n-butanol: Glacial Acetic Acid: Water (8:2:2) (v/v/v) solvent system. UV and IR spectra of isolated and standard marker compound (Levodopa) showed similar results. Chemical test of Levodopa showed the positive result. All above observation and results confirm the isolated compound was Levodopa

REFERENCES

1. Kokate CK, Gokhale SB, Purohit AP, 2002. Pharmacognosy. 23rded. Pune: Nirali Prakashan. 35, Pages 119-121.
2. Nainwal P, 2010. Pharmacovigilance of herbal medicines: an intangible approach. IJPSR. 1(1), Pages 60-65
3. Mosihuzzaman M, Choudhari M, 2008. Protocols on safety, efficacy, Standardization and documentation of herbal medicine (IUPAC technical report). Pure Appl Chem. 80(10), Pages 2195-2230.
4. Mali D, 2011. Dissertation report on standardization of herbal medicines: Triphala Churna. Shivaji University, Maharashtra.

5. Sahoo N, 2010. Herbal drugs: Standards and regulation. Fitoterapia. 30(60).
6. Sharma A, 2008. Herbal Medicine for Market Potential in India: An Overview. Academic Journal of Plant Sciences. 1(2), Pages 26-36.