



Role of cytokines in pathogenesis of rheumatoid arthritis (chalcone)

Hem Raj Vashist*, Avneet Gupta, Neeraj Upmanyu

Department of pharmacy CMJ University Jorabat, Rai Bhoi, Meghalaya, India

Corresponding author: Hem Raj Vashist, Shimla_pharmacy@rediff.com,

Department of pharmacy CMJ University Jorabat, Rai Bhoi, Meghalaya, India

© The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by-nc/4.0/>). See <https://jmpas.com/reprints-and-permissions> for full terms and conditions.

Received – 11 April 2015, Revised – 22 May 2015, Accepted – 20 June 2016 (DD-MM-YYYY)

[Refer This Article](#)

Hem Raj Vashist, Avneet Gupta, Neeraj upmanyu, 2016. Role of cytokines in pathogenesis of rheumatoid arthritis. Journal of medical pharmaceutical and allied sciences, V 5 - I 4, Pages -351 – 355. Doi: <https://doi.org/10.55522/jmpas.V5I4.0092>.

ABSTRACT

Cytokines are polypeptides autacoids produced by phagocytes and some other cells. A large number of cytokines are active in the joints of patients with rheumatoid arthritis (RA). It has been evidently proved that cytokines play fundamental role in bone reabsorption and connective tissue destruction and also in complications associated with RA. By knowing the sinister role of cytokines to induce rheumatoid arthritis in patient, the present study is focused on the review of different cytokines which involve in the pathogenesis of the disease. From the literature available from different article on cytokines it is clear that cytokines like Tumor necrosis factor (TNF- α), chemokines and Interleukins (IL) are responsible for the pathogenesis of rheumatoid arthritis. From the previous literature it was concluded that Cytokines are responsible for the pathogenesis of rheumatoid arthritis. Safe intervention can be evaluated by understanding this fact.

Keywords: Cytokines, Polypeptides, Autacoids, Rheumatoid Arthritis, Bone Reabsorption, Tissue Destruction.

INTRODUCTION

(Greek *cyto-*, cell; and *-kinos*, movement) Cytokines are known to be pivotal mediators in the RA process. Cytokinins are polypeptide substances produced by activated lymphocytes (lymphokines) and activated monocytes (monokines) may act on self cell producing them or on other cells. These polypeptides are known to mediate some aspect or other of the inflammatory process. The effector phase of both natural and acquired immunity are in large part mediated by cytokines. Cytokines are diverse meaning and are not alike. Different types of cytokines are produced by cells like i) colony stimulating factor which stimulate production of blood cells ii) Growth and differentiation factor which function primarily in development and immunoregulatory and proinflammatory cytokines. Cytokines intera viz interleukins, interferon and TNF- α that function in immune system. Cytokines interact with cells of the immune system in order to regulate the body's response to disease and infection. Cytokines also mediate normal cellular processes in the body. Cytokines serve as molecular messenger between cells, with regard to arthritis, cytokines regulate various inflammatory responses. Overproduction or inappropriate production of certain cytokines by the body result in arthritis. For example, it has been found that interleukin-1


(IL-1) and tumor necrosis factor-alpha (TNF- α) are produced in excess in [rheumatoid arthritis](#) where they are involved in inflammation and tissue destruction. Cytokines act in the response of injury where it helps to exert innate immune response. The innate immune system which act as double edged sword for the host. Many pro-inflammatory cytokines including TNF- α , chemokines and growth factors are expressed in diseased joints. Rheumatoid arthritis joint synovial membrane infiltration and cartilage degeneration whereas a normal joint show a normal cartilage and proper shape of synovial membrane. Proinflammatory cytokines like TNF- α and IL-1 β are meant mostly to be involved in the rheumatoid arthritis. Single cytokine is alone able to exert its pleiotropic activity and different cytokines combinely shows redundancy like TNF- α alone can induce inflammation, fever, sleep anorexia and alteration in neuroendocrine system, whereas the overproduction of TNF- α , IL-1, IL-6, IL-17, IL-18 and so many other interleukins shows their action to induce RA. The Fig1&2 depicted below are showing a normal knee joint and arthritic joint. Fig1&2.

Proinflammatory cytokinins like TNF- α , IL-1, IL-6, GM-CSF and chemokines like IL-8 play a pivotal role in the pathogenesis of


Rheumatoid arthritis. It is now accepted that TNF- α and IL-1 play a major role in RA. Cytokines are a group of chemicals like Tumor necrosis factors (TNF) and Interleukins (IL). The member of tumor necrosis factor involve TNF- α , TNF- β and TNF- γ whereas interleukines involve IL-1, IL-2, IL-6 and IL-23 and many more. The most involving cytokines in the RA are TNF- α , IL-1, IL-6 and IL-2. Several researchers have reported the role of these cytokines in the pathogenesis of RA. Present study is therefore focused on the review on the role of cytokines (TNF- α and IL-1 to IL-10) in RA [1].

Material TumorNecrosisFactor (TNF)

TNF- α was identified in 1975 as the factor in serum isolated from endotoxin treated mice. It is a pleiotropic, pro-inflammatory, pro-apoptotic cytokine that regulates the activity of neutrophils, eosinophils, and T & B lymphocytes and modulates the properties of the vascular endothelium.



Tumor necrosis factor (TNF), cachexin or cachectin formerly known as tumor necrosis factor alpha or TNF- α is a cytokine involved in systemic inflammation and is a member of a group of cytokines that stimulate the acute phase reaction. It is produced chiefly by activated macrophages, although it can be produced by other cell types as well. The primary role of TNF is in the regulation of immune cells.



Being an endogenous pyrogen, TNF is able to induce fever, induce inflammation, to induce apoptotic cell death, to induce cachexia, to induce sepsis (through IL1 & IL6 production), and to inhibit tumorigenesis and viral replication. The primary role of TNF is in the regulation of immune cells. It has been found that dysregulation of TNF production leads to induce variety of human diseases, including Alzheimer's disease, inflammatory bowel disease (IBD) cancer and major depression.

Role in pathogenesis of arthritis. The actions of TNF- α is important in the pathogenesis of RA include its ability to induce the production of other proinflammatory cytokines, including IL-1 and IL-6, together with its ability to induce the production and release of chemokines. TNF- α is a pleiotropic, proinflammatory mediator which is produced in response to infection to confer immunity to the host. It is now accepted that TNF and IL-1 produced locally in the inflamed synovial joint contribute directly or indirectly to the pathogenesis of rheumatoid arthritis (RA). The effect of the TNF- α in infection is beneficial but tight regulation of TNF- α production is required to protect the host from its detrimenta [2].

Activities. Deregulated production of TNF- α is thought to play pivotal role in pathogenesis of RA. T-cells are found responsible for the secretion of TNF by direct contact mediated interaction. Histological studies of synovium in RA have indicated that this tissue is very cellular and that several different cell type including macrophages and T-cells are in close proximity. Direct contact mediated interaction using transformed T- cells and monocytes have been found to play a major role in inducing the release of TNF- α , IL10 and metalloproteinase specific surface interaction molecule that are reported to mediate induction of monocyte cytokine synthesis includes CD69, IFA1 CD44 CD40 membrane TNF[29] and signaling lymphocytic activation molecule equilibrium induced by cytokine stimulated T-cell (Tck), monocytes(MQ),monocytes(Tcr) cell receptor dependent stimulated T cells.

Evidences for the role of TNF- α in the pathogenesis of RA has been reported by showing its presence in synovial fluid. The role of TNF- α in early joint swelling in RA has found.

The presence of TNF- α in rheumatoid arthritis serum and synovial fluid has been presented by using several experimental models for rheumatoid arthritis. Significant concentrations of TNF- α and IL-1 has been reported in arthritic joints of experimental antigen induced arthritic rabbit.

In one of the study TNF- α has been reported to be secreted by activated mast cells. Mast cells are equipped with FC receptors for binding to autoantibodies and the receptor for binding to complement usually in the form of antibody compliment complex. Mast cell produces large amount of TNF- α and IL-1 cytokines that play a critical role in perpetuating the inflammation. Mast cell might have received the signal when bound by autoantibodies and complement fragments and release TNF- α and IL-1,

contributing to the pathology in the synovium that result in erosive arthritis.

Mechanism of TNF- α to lead inflammation to RA has explained by one more way as TNF- α is responsible to trigger the release of other cytokines like proinflammatory cytokines (IL-1, IL-6, IL-23 and GM) together with its ability to induce the production and release of chemokines (IL-8) that attract leukocytes from the blood into inflamed tissue. This process is facilitated by upregulation of key integrins and adhesion molecules on endothelium which finally causes the destruction of proteolytic and metalloproteinase enzyme which bring about the destruction of subchondrial boneDropwise spectrometer. The chemical shift were recorded

Ar-CO-CH=CH-Ar' in parts per million (ppm) and were reported relative to the TMS. Mass spectra data were recorded on Waters Q-TOF Premier-HAB213 Scheme-01 [3].

Adopting conventional Claisen-Schmidt procedure parallelly in our lab we encountered problems such as prolong reaction time, excessive solvent requirement, sticky product system in ESI mode. The FT-IR spectra of synthesized compounds were recorded on PerkinElmer Spectrum version 10.03.06. The melting point was recorded by open capillary method and is uncorrected [4].

Overproduction of TNF- α is regulated by using TNF- α inhibitors and also called TNF- blockers. These blockers bind to TNF- α and prevent TNF- α attaching to cell surface receptors. Enbrel (Etanercept), remicade (Infliximab) and humiradalimumab are TNF blockers.

Interlukines-(inter-in;lukine-leukocytes). Interlukines are the member of cytokines which are meant play a major role in the pathogenesis of RA. The role of different interlukines in the pathogenesis of RA is discussed here one by one.

Interleukin -1(IL-1)- Role of IL -1 cytokine in the pathogenesis of RA has been implicated by significant experimental evidences. To understand the role of cytokines in RA the different cells involved should be discussed. The pathogenesis of RA depends upon a number of cell types (Fig- 4). Macrophage after activation express HLA- Class II molecule. Because of their proximity to T-cell they may function as antigen presenting cell and perpetuate immunological response within the joint. Chondrocytes get involved in these properties and have the ability to produce cartilage specific antigen. In RA chondrocytes release degradative enzyme and reduces its capacity to synthesize new matrix components. T-cells in the rheumatoid joints have a memory CD4+ phenotype which express various activation markers and are found in close proximity to antigen presenting cells[. β -Cells are also found in large number in the rheumatoid joints β -Cells are responsible to produce immunoglobulines and autoantibodies i.e. rheumatoids factor and anticholagen antibodies which form immune complexes that can local inflammation as well as IL-1and TNF- α released by macrophage.

Another type of cell which plays a role in pathogenesis of RA is synoviocytes and endothelial cells. These cell express specific cell adhesion molecule (CAM) on their surface along with intracellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule 1 (VCAM-1) and E-selectin which control the infiltration of inflammatory cells into the joint. The fibroblast like synoviocytes predominates into the pannus IL-1 and TNF- α upregulates the expression of CAMs on endothelial cell their by directing emigration of blood cells from circulation into the synovium.

IL-1 Family consist of three members viz IL-1 α , IL1 β produce the biological effects attributed to the cytokines and IL-1 receptor antagonist(IL- 1Ra) is an endogenous inhibitor that block the action of the other two members. The three dimensional structure of the family members are related. Each member in the family has a distinct amino acid sequence. IL-1 α , and IL-1 β are produced as 31kDa precursor protein and termed as pro IL-1 α and pro IL-1 β . Both IL-1 α is generally retained within the cell or expressed on the cell surface which is believed to act as autocrine messenger. IL-1 β is secreted then and act on other cell to produce its biological action. IL-Ra is synthesized and secreted as 17kDa proteins by the same cell. Each members of IL-1 binds with high affinity to specific receptor located on the surface of the target cell. Binding of IL-1 α or IL-1 β to type 1 IL-1 receptor (IL-1R1) which is enhanced by an accessory protein IL-1R- AcP leads to intracellular signals transduction and regulation of gene expression and as a result cellular responses begining to the type II IL-1 receptor (IL-1RII) on these cells does not cause cell activation because the receptor contain very short cytoplasmic domain that is unable to signal transducer signal following IL-binding. Therefore IL-1R-II receptor acts as decoy receptor. IL-1Ra binds to both IL-1R1 and IL-1RII and does not induce signal transduction. Instead, it competitively antagonize the binding of IL-1 α and IL-1 β and thereby reducing the effect of these cy.

Interlukine-2(IL-2) - IL-2 T-cell growth factors produced by activated helper, is an important immunoregulatory molecule required for normal lymphocyte function. The synthesis of IL-2 leads to production of interferon gama (IFN- γ) by T-cell. The synthesis of IL-2 is generally consider as a proinflammatory cytokine that exacerbate Th-1 mediated disease states such as autoimmune arthritis. A growing body of evidence from both animal model of arthritis and human rheumatoid arthritis suggest that Th-2 type cytokines such as IL-4 and IL-10 can protect against arthritis whereas Th-1 type cytokine such as IL-2 and IFN- γ can be proinflammatory. Number of observations support for this concept. The RA synovium is infiltrated with CD4+T cell Th-1 phenotype. In murine collagen induced arthritis (CIA) a model of RA Abs can inhibit disease onset. It is surprising that most studies in cytokine has failed to detect IL-1 [3].

protein, in RA synovial fluid. Few evidences has been proved experimentally for the presence of IL-2. The failure of detection of IL-2 may relate to the timing of synovial sampling. In a study the serial sampling of synovium of RA patient was avoided because of difficulty. The experiment was tested on murine collagen-induced arthritis model to analyze mRNA cytokine pattern at various stages of autoimmune arthritis. These studies demonstrated that IL-2 mRNA decline during the chronic stage of the disease. Further support for this fact comes from the record demonstrated that IL-2 protein is significantly increased in peripheral blood mononuclear cells from patient with early but not late RA. In order to characterize T-lymphocyte function in RA the production of IL-2 employing T-cells from synovial fluid(SF)Synovial tissue and peripheral blood(PB) of patient with RA. Total nineteen patient 15 female and 4 male with definite or classical RA were studied. SF was obtained from 15 and ST from IL-2 assay and IL-2 activity was detected using murine IL-2 dependent cytotoxic T-cell line (CTLI-2).

Interleukin-3 (IL-3). The synthesis of IL-3 is regulated by CD4+ cell. An amount of IL-3 has been reported in synovial fluid by using CIA mice model. IL-3 is Known to be a potent growth promoting cytokine and is found to associated with neurologic disorders. IL-3 primarily produced by activated T-cells stimulates growth and differentiation of monocytes, basophils and other leukocytes population from the bone marrow in an immune response. It has been demonstrated that IL-3 plays an important role in the onset of RA, an autoimmune disease. Along with its growth regulatory action IL-3 also induce the release of IL-6 and IL-4. But the release of IL-4 is 3 time less than IL-6. This fact has proved in CIA induced arthritis in mice. In the same model an amount of IL-3 was found. The recombinant IL-3 induce histamine release from human basophiles also leads to some aggressive form of inflammation.

Interleukin-4(IL-4). Is an anti-inflammatory cytokine and help to control proinflammatory cytokines. IL-4 is a member of CD132 dependent cytokine which also comprise IL-2, 7, 9, 15 and 21. IL-4 have two receptors Type I which is composed of IL-4R α and CD132(IL-2 R γ or γ c) and type II receptor composed of IL-4R α and IL-13R α chain [64]. IL-4 has largely been associated with anti-inflammatory since it decreases monocytes and macrophages functions and induces Th2 type responses. IL-4 also has been reported to possess proinflammatory functions such as recruitment of inflammatory cells on the endothelial cell surface and increasing eosinophils, macrophages and B cell chemoattraction. This dual nature of IL-4 confirm its anti and proinflammatory activity. Several members of the CD132-dependent cytokines include IL-2, IL-4, IL-7, IL-9 and IL-21 are involved in diverse inflammatory disorder among these IL-4 is best known for its anti- inflammatory activity. IL-4 has found to induce IL-1 receptor antagonist in neutrophils. In one more study by using the murine ear air pouch model it has been reported that IL-4 possess proinflammatory activity invivo based on ability to attract leukocyte

including neutrophils. By analyzing two polymorphs of IL-4 and polymorph of IL-4RA genes in patient with RA, the role of IL-4 in this disease has been reported.

Interlukin-5(IL-5). Is produced by T-helper 2 cells which stimulate B-cell growth and increase immunoglobulin secretion. IL-5 Th2 cytokine that is a part of homopoetic family. Being a hematopoietic cytokine, IL-5 plays a key role in the differentiation, maturation, and survival of eosinophils derived from bone marrow precursors. The link between IL-5 and eosinophils is highly conserved in mammals which suggest a selective pressure for maintenance of this function in the immune repertoire. In one of the study mouse strains with genetically altered IL-5 expression have been used to understand what changes occur to the immune system as a result of its dysregulation. Depending on the transgenic construct, IL-5 overexpression has been shown to cause moderate to massive eosinophilia, lower bowel inflammation, and enhanced autoantibody production. IL-5 has been shown to be essential for eosinophils maturation and survival. Over all the action of IL-5 by activating Th1 receptor shows its anti-inflammatory action. But its role in the production of auto-antibody also shows its role in autoimmune disease [4].

Interlukines-6(IL-6). IL-6 was identified in 1986 as a B cell regulatory factor and is now recognized as mediating as pleiotropic functions which include effect on the maturation and activation of B and T cells, macrophages, osteoclasts, chondrocytes and endothelial cells and broad effect on haematopoiesis in the bone marrow. IL-6 is produced by a variety of cell types in response to infection, trauma, and immunologic challenge. IL-6 plays a prominent role in disease processes and has both proinflammatory and anti- inflammatory characteristics. It promotes inflammatory events through the expansion and activation of T cells and the differentiation of B cells. IL-6 has also demonstrated a protective role in disease processes. For example, in a septic shock model, IL-6 protected mice against death by suppressing acute neutrophils accumulation caused by intratracheal administration of endotoxin. IL-6 produces many other effects throughout the human body, both locally and systemically. IL-6 is relevant to several disease such as diabetes, Atherosclerosis, depression, Alzheimer disease [, systemic lupus erythematosus and Rheumatoid arthritis.

Interleukin 6 (IL-6) plays a pivotal role in the pathophysiology of rheumatoid arthritis (RA). It is found in abundance in the synovial fluid and serum of patients with RA and the level correlates with the disease activity and joint destruction. IL-6 can promote synovitis and joint destruction by stimulating neutrophils migration, osteoclasts maturation and vascular endothelial growth factor (VEGF)-stimulated pannus proliferation. Concentrations of IL-6 have been reported in synovial fluid and serum/plasma from patients with rheumatoid arthritis.

Several studies have confirmed that excessive amounts of IL-6 and

soluble IL-6R are produced in the joints of patients with RA, particularly the synovium, which is a thin tissue layer covering the joint from the inside. These increases contribute to inflammation, swelling, joint damage. **Interlukin-9(IL-9)** Interleukin-9 is a cytokine produced by Th2 cells. IL-9 was first identified as a mouse T cell growth factor [99] but using in vitro assay systems it has also been demonstrated to modulate B cell maturation and to lymphopromote proliferation and differentiation of mast cells and hematopoietic progenitors. IL-9 was originally described as a mast cell growth factor for its ability to enhance the survival of primary mast cells and to induce their production of the inflammatory cytokine IL-6 thus indirectly plays a role in the progress of RA. IL-9 also stimulates the production of mast cell proteases and the high-affinity IgE receptor, suggesting that IL-9 primes mast cells to respond to allergen via increased cell surface expression of the high-affinity IgE receptor and the production of inflammatory mediators, including IL-6 and several proteases [5].

Interleukin-10 (IL-10) Interleukin-10 functions as an effective immunomodulatory molecule based on two important functions viz. inhibition of cytokine synthesis and down regulation of antigen presenting cells. IL-10 has been described as murine Th2 cell product. Role of IL-10 in arthritis has been reported by taking the peripheral blood sample from RA patient and from synovial joint fluid. The anti-arthritis activity of IL-10 has been given experimentally by using human and CIA mice model with the observation that anti-IL-10 helps in the enhancement of arthritic condition more in both models whereas injected exogenous IL-10 lower the severity of the disease. The action of IL-10 is correlated with that of IL-4. Instead of the anti-inflammatory activity IL-10 are also

meant to release some proinflammatory cytokines. age and destruction associated with RA because of IL-6 [6].

REFERENCES

1. Parry SL, Sebbag M, Feldmann M, Brennan FM, 1997. Contact with T cells modulates monocyte IL-10 production: role of T cell membrane TNF- alpha. *J Immunol.* 1(58), Pages 3673-3681.
2. Wu J.Z Cheng, C C Shen, L.L Wang, antioxidant ability on H2O2-induced apoptosis in PC12 cells. *Int. J. Mol. Sci.*, Pages 18525–18539.
3. Ford ES, Giles WH, Dietz WH, 2002. Prevalence of metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA.* 287(3), Pages 356–359. Doi:10.1001/jama.287.3.356.
4. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (May 2001). "Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III)". *JAMA: the Journal of the American Medical Association.* 285(19): Pages 2486–97. Doi: 10.1001/jama.285.19.2486.
5. Wim B van den Berg, Leo A B Joosten, George Kollias, Fons A J, 1999. van de Loo Role of tumour necrosis factor α in experimental arthritis: separate activity of interleukin 1 α in chronicity and cartilage destruction. *Ann Rheum Dis.* 58(I), Pages I40–I48.
6. Benjamin Yang., 2002. Mast cells make and release TNF-alpha in synovium, play pronounced role in arthritis. *Discovery Medicine.* 1(11), Page 2.