



## Research article

**Implication of heat shock proteins in periodontal disease**

Ashank Mishra\*

Department of Periodontics, Sri Sai College of Dental Surgery, Vikarabad, India

**Corresponding author:** Ashank Mishra, ✉ [Vijdeep@gmail.com](mailto:Vijdeep@gmail.com),  
Department of Periodontics, Sri Sai College of Dental Surgery, Vikarabad, India

© The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by-nc/4.0/>). See <https://jmpas.com/reprints-and-permissions> for full terms and conditions.

**Received** – 20 August 2016, **Revised** - 25 September 2016, **Accepted** – 23 October 2016 (DD-MM-YYYY)

[Refer This Article](#)

Ashank Mishra, 2016. Implication of heat shock proteins in periodontal disease. Journal of medical pharmaceutical and allied sciences, V 5 - I 5, Pages -374 – 375. Doi: <https://doi.org/10.55522/jmpas.V5I5.0097>.

**ABSTRACT**

Heat shock proteins are found in virtually all living organisms, from bacteria to humans. They stabilize proteins and are involved in the folding of denatured proteins. During the infection, HSP from several bacterial species are recognized by host as immunodominant antigens. Periodontopathic bacteria have sequence homology with human HSP 60 at amino acid level, which raises the possibility of cross reactivity. The present review gives an insight into the role of HSPs in the etiopathogenesis of periodontal disease

**Keywords:** Heat shock proteins, Periodontopathic bacteria, cross reactivity.

**INTRODUCTION**

Heat shock proteins (HSP) are a class of functionally related proteins whose expression is increased when cells are exposed to elevated temperatures or other stress. This increase in expression is transcriptionally regulated. The dramatic upregulation of the heat shock proteins is a key part of the heat shock response and is induced primarily by heat shock factor (HSF). HSPs are found in virtually all living organisms, from bacteria to humans [1].

Heat-shock proteins are named according to their molecular weight. For example, Hsp60, Hsp70 and Hsp90 (the most widely-studied HSPs) refer to families of heat shock proteins on the order 60, 70 and 90 kilodaltons in size respectively. The small 8 kilodalton protein ubiquitin, which marks proteins for degradation, also has features of a heat shock protein [2].

Heat shock proteins are present in cells under normal conditions, but are expressed at high levels when exposed to a sudden temperature jump or other stress. Heat shock proteins stabilize proteins and are involved in the folding of denatured proteins. High temperature and other stresses such as altered pH and oxygen deprivation, make it more difficult for proteins to form their proper structures and cause some already structured proteins to unfold. Left uncorrected, misfolded proteins form aggregated that may eventually kill the cell. Most

heat shock proteins are molecular chaperones. Chaperones aid in the transport of proteins throughout the cell's various compartments [3].

Under normal conditions, heat shock proteins are required for cellular metabolism and help newly synthesized poly peptides fold, thus preventing premature interactions with other proteins.

In 1962, Ritossa reported that heat and the metabolic inhibitor dinitrophenol induced a characteristic pattern of puffing in the chromosomes of *Drosophila*. This discovery eventually led to the identification of the heat-shock protein (HSP) or stress proteins whose expression these puffs represented. Increased synthesis of selected proteins in *Drosophila* cells following stresses such as heat shock was first reported in 1974 [4].

**Classification of heat shock proteins**

The principal heat-shock proteins that have chaperone activity belong to five conserved classes; HSP33, HSP60, HSP70, HSP90, HSP100 and the small heat-shock proteins (sHSPs).

**Extracellular stress proteins*****Osteonectin/ (Secreted Protein Acidic and Rich in Cysteine) SPARC***

Osteonectin/SPARC is a 32 kDa phosphorylated glycoprotein originally purified from bone tissues, has been identified in several tissues, including placenta, platelets and

fibroblasts. Analyses of protein structures have indicated that osteonectin and SPARC are identical [5].

### **Thrombospondin**

Thrombospondin (TSP) is a trimeric extracellular

It is interesting that mRNA of two metalloproteinases, collagenase and stromelysin which are produced in large quantities in rheumatoid arthritis, diabetes, periodontal disease and connective tissue undergoing remodeling are expressed by heat shock.

### **Role of HSP in periodontal diseases**

HSP are perhaps, are investigated extensively as a self antigens in periodontal diseases. On exposure to environmental factors, these proteins participate in various processes like folding and translocation of polypeptides across the membranes. These HSP can also be referred as molecular chaperones.

During the infection, HSP from several bacterial species are recognized by host as immunodominant antigens. Periodontopathic bacteria have sequence homology with human HSP 60 at amino acid level. These proteins are found in the tissues samples of periodontitis lesions [6].

HSP 60 is considered as very immunogenic and the autoimmune response to it could be touted as the initiator of inflammatory diseases. It is hypothesized that the periodontopathic bacteria stimulate the cells of periodontium to secrete amount of HSP 60 expression which in turn leads to exaggerated secretion of proinflammatory cytokines by macrophages. (Pleguezuelos et al)<sup>6</sup>

It was reported initially that the expression of HSP 60 was higher in gingivitis and periodontitis tissue extracts compared to healthy tissue. (Lundqvist et al 7, Petit et al 8, Ueki et al 9) Yamazaki et al 10 and Choi et al 11 demonstrated HSP specific T cell might be involved in the immunopathogenesis of periodontal diseases.

Recent paper by Nethravathy et al 12 reported increased levels of Circulatory HSP 60 in periodontal disease compared to health individuals.

Molecular mimicry: Role of HSP 60 in periodontal diseases- cardiovascular disease link 13.

The concept of molecular mimicry was introduced by Raymond Damian in 1964. He suggested that antigenic determinants of microorganisms may share homology with antigenic determinants of their hosts. It was hypothesized that molecular mimicry may induce cross reactive immune response with host antigens, resulting in an autoimmune disease like mechanism.

This mechanism was touted as fundamental for atherogenesis. Endothelial cells under stressful conditions express heat shock proteins. In addition, periodontopathic bacteria i.e. *P.gingivalis* express heat shock proteins like proteins such as

GroEL. It was proposed that structural homology exists between HSP 60 and GroEL of *P.gingivalis*, the anti *P.gingivalis* immune response

cross-reacts with human HSP 60 expressed on endothelial cells, resulting in endothelial dysfunction and atherosclerosis [7].

### **CONCLUSION**

Heat shock proteins and their role in etiopathogenesis of periodontal disease would aid in diagnosing and treating the disease. HSP could be considered as a potential candidate antigen for periodontal vaccine.

### **REFERENCES**

1. Csermely P, Yahara I, 2003. Heat Shock Proteins. In: Keri G, Toth I, editors. Molecular Pathomechanisms and New Trends in Drug Research CRC Press. Pages 67-75.
2. Lundqvist C, Baranov V, Teglund S, 1994. Hammarstrom ML Cytokine profile and ultrastructure of intraepithelial gamma delta T cells in chronically inflamed gingiva suggest a cytotoxic effector function J Immunol. 153, Pages 2302-2312.
3. Manyahi J, Matee MI, Majigo M, 2014. Predominance of multi-drug resistant bacterial pathogens causing surgical site infections in Muhimbili National Hospital, Tanzania. BMC Res Notes. 7, Pages 500. Doi: 10.1186/1756-0500-7-500.
4. Muhammad UK, Adamu TM, Binji Z, 2014. Prevalence of  $\beta$ -lactamase production among pathogenic bacteria isolated from surgical site and wound infection among patients admitted in some selected hospitals in Sokoto Metropolis, Nigeria. Int J Env. 3(3), Pages 104-112.
5. Poirel L, Walsh TR, Cuvillier V, 2011. Multiplex PCR for detection of acquired carbapenemase genes. Diagn Microbiol Infect Dis. 70(1), Pages 119-123.
6. Sharif L, Obaidat M, Al-Dalalah M, 2013. Food Hygiene Knowledge, Attitudes and Practices of the Food Handlers in the Military Hospitals. FNS. Doi: <http://dx.doi.org/10.4236/fns.2013.43033>. J
7. Petit MD, Wassenaar A, van der Velden U, 1999. Loss of Depressed responsiveness of peripheral blood mononuclear cells to heat shock proteins in periodontitis patients J Dent Res. 78, Pages 1393-1400.