



Research article

Isolation and Characterization of a Dihydrochalcone from *Detarium senegalense* (J. F. Gmel.) Stem Bark and Biological Activity of Different Fractions

Abdel Karim M*, A F Sakina

Department of Chemistry, Sudan University of Science and Technology, Khartoum, Sudan

Corresponding author: Basak Silpi, ✉ Vijdeep@gmail.com,

Department of Pedodontics and Preventive Dentistry, Sharad Pawar Dental College, Datta Meghe Institute of Medical Sciences (Deemed University), Maharashtra, India

© The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by-nc/4.0/>). See <https://jmpas.com/reprints-and-permissions> for full terms and conditions.

Received – 20 August 2016, Revised - 25 September 2016, Accepted – 23 October 2016 (DD-MM-YYYY)

Refer This Article

Abdel Karim M, A F Sakina, 2016. Isolation and Characterization of a Dihydrochalcone from *Detarium senegalense* (J. F. Gmel.) Stem Bark and Biological Activity of Different Fractions. *Journal of Medical, Pharmaceutical, and Allied Sciences*, V 5 - I 5, Pages -382 – 385. Doi: <https://doi.org/10.55522/jmpas.V5I5.0100>.

ABSTRACT

Information on the constituents of medicinal plants used in Sudanese traditional medicine is very scarce. Hence, this study was set to investigate the flavonoids of *Detarium senegalense* which is widely used by local healers to treat an array of human disorders. Phytochemical screening of *Detarium senegalense* ethanolic extract revealed the presence of sterols, triterpenes, flavonoids, tannins, saponins, anthraquinones and cyanogenic glycosides. A dihydrochalcone was isolated from stem bark and its structure was partially elucidated on the basis of its spectral data (IR, UV, ¹H NMR and MS). The isolate was evaluated, *in vitro*, for its antimicrobial activity against six standard human pathogens: two Gram positive (*Staphylococcus aureus* and *Bacillus subtilis*), two Gram negative (*Pseudomonas aeruginosa* and *Escherichia coli*) bacteria and two fungal species (*Aspergillus niger*, *Candida albicans*) and promising results suggested that the dihydrochalcone is a plausible lead for further optimization.

Keywords: *Detarium Senegalense*, Isolation, Dihydrochalcone, Antimicrobial Activity.

INTRODUCTION

In the family Fabaceae, *Detarium*, is a genus being represented by 8 species. In west African forests, only three species are found: *Detarium macrocarpum* Harms, *Detarium microcarpum* Guillemin and Perrottet and *Detarium senegalense* J.F. Gmelin. These species are of pharmacological interest ^[1].

Detarium senegalense J. F. Gmel. or Sweet detar is a species of plant in the Fabaceae family, subfamily: Caesalpinioideae and tribe: Detarieae. (Also placed in Leguminosae and Caesalpinieae). It grows naturally in the drier regions of west and central Africa extending from Senegal and Gambia east to north-east tropical Africa and Sudan. *Detarium senegalense* fruit seems to be a very healthful nutrient. It contains (Per 100 g): 27 mg calcium,

48 mg phosphate, 0.14 mg thiamin, 0.05 mg riboflavin and 1000–2000 mg ascorbic acid. Fruit pulp contains (per 100g): 2.8 mg iron, 0.6 mg niacin, and about 1200 mg vitamin C. Lesser quantities of other vitamins and minerals were reported. Seeds are claimed to contain

12% protein, and are rich in lysine-the rare amino acid- and tryptophan. Stem bark and seeds were found to be rich in vitamins comprising (mg/100g) riboflavin 0.62 – 0.60, thiamin 0.14 – 0.27, niacin 2.06 – 8.11 and ascorbic acid 83.60^[9]. The nutritional value of *Detarium senegalense* seeds seems to justify its use in the traditional treatment of skin diseases.

Phytochemical studies of the seeds and stem bark revealed the presence of: alkaloids, flavonoids, tannins, phenols and saponins. *Detarium senegalense* seeds contain a large amount of water-soluble, non-starch polysaccharide (s-NSP) suggesting important nutritional value. The main monosaccharide residues are: glucose, xylose, and galactose.

The fruit pulp is edible, eaten raw or cooked. The seed is oily and edible, and pounded seed is used as cattle feed. Seed flour is used traditionally in Nigeria as an emulsifying, flavouring and thickening agent in foods. It is also employed as a soup thickener.

GC-MS analysis of the petroleum ether seed extract revealed the presence of ten constituents being dominated by oleic and linoleic acids (30.8 and 44.1% respectively). Hanan demonstrated that the petroleum ether fraction exhibited significant antibacterial activity against a panel of human pathogens. Sowemimo *et al*, claimed antifungal activity against *Aspergillus flavus*, *Aspergillus niger* and *Penicillium notatum*.

A xyloglucan (a non - starch polysaccharide) extracted from *Detarium senegalense*, has considerable promise in the treatment of diabetes and hyperlipidaemia. The rheological and other properties of this component indicate a considerable commercial potential in the food, drugs and chemical industries. Recently There was considerable interest in the functionality of dietary polysaccharides starch and non-starch polysaccharides (NSP) with respect to prevention and treatment of disease such as diabetes, coronary heart disease, and arthritis [1].

Detarium senegalense is widely used in ethno medicine to treat an array of human disorders. Bark is claimed to treat dropsy, swelling and oedema. A decoction of bark is used to treat digestive disorders, bronchitis, pneumonia and stomach –ache, while bark powder is applied to wounds and burns. Bark pulp is eaten as a general tonic and in treatment of tuberculosis. The bark is also used for headache, back-pain, sore throat and painful menstruation and aqueous bark extract is used to cure indigestion. Root decoctions are administered as anodyne and to treat intestinal disorders, convulsions and anaemia. Leaf decoction is said to treat fever, trypanosomiasis, dysentery, anaemia, conjunctivitis, arthritis, inflammations, fractures, boils and skin infections. Fruit pulp is a treatment for kidney troubles, spinal tuberculosis, syphilis, cough, rheumatism and leprosy. In one study a *Detarium senegalense* meal was shown to elicit significant reduction in plasma glucose levels in studied human subjects. Seeds have been effective in controlling blood glucose levels in diabetic individuals. Also, they are taken as antidote against snake bites and arrow poison. Seeds are also used by local healers as emetic. Bark and fruit are used to treat leprosy and pulmonary troubles. Bark and leaf generally heal skin and mucosa. Leaves used traditionally as wash for itch, enema for dysentery and eye wash for conjunctivitis. Root and fruit are used as painkillers. Root and leaves decoctions are used in paralysis, meningitis and difficult delivery. *In vivo* studies testified promising antidiabetic effect for gum. Seeds, leaves and roots decoctions are used as remedy for venereal diseases, intestinal worms, diarrhea, urogenital infections, hemorrhoids, rheumatism, stomach ache, leprosy and malaria.

Leaves from the trees have demonstrated antiviral activity against a number of human and animal viruses and the bark has shown antibacterial activity against many pathogenic bacteria, justifying the medicinal properties of the plant. An anthocyanin alkaloid 2-

methoxyamine 3,4,5,7-tetrahydroxyanthocyanidine has been isolated from the stem bark. *In vitro* antibacterial studies showed that the isolated compound successfully inhibited a panel of human pathogens. Such results authenticate the use of *Detarium senegalense* in phytomedicine for the treatment of infections.

In continuation of our interest in the constituents of medicinal plants used in Sudanese traditional medicine, this study was designed to investigate the constituents of the Sudanese material of *Detarium senegalense* and to evaluate the antimicrobial activity of its fractions, hoping to establish a rationale for its ethno-medical uses [2].

MATERIALS AND METHODS

Plant Material

The stem barks of *Detarium senegalense* were collected from western Sudan. The plant was authenticated by direct comparison with a herbarium sample [3].

Solvents

Analytical grade solvents were used. Methanol - HPLC grade- was used for spectroscopic purposes (Loba, India).

Equipments

The ultraviolet lamp used in visualizing TLC plates and paper chromatography was a multiband UV λ_{max} (254 / 365 nm) portable ultraviolet lamp, a product of Hanovia Lamps (6 watt S/Y and L/W). Ultraviolet absorption spectra were obtained in spectroscopic methanol on UV - Visible Spectrophotometer (Shimadzu).

Infrared spectra were generally obtained in potassium bromide (KBr) discs using Perkin- Elmer, FTIR, model 1600-Jasco. The electron impact ionization (EIMS) mass spectra were obtained on a solid probe using Shimadzu QP- class-500.

¹HNMR spectra were obtained on a Bruker AM 500 spectrophotometers (Germany) operating at 500 MHz in spectroscopic grade DMSO-d₆. The chemical shifts values are expressed in δ (ppm) units using (TMS) as an internal standard and the coupling constants (J) are expressed in Hertz (Hz).

Methods Stepwise procedure for use of shift reagents for UV

The UV spectrum of the compound in methanol was first recorded.

3 drops of NaOMe reagent were added to the sample and the NaOMe spectrum was recorded, and after 8 minutes the NaOMe spectrum was re- recorded.

6 drops of AlCl₃ reagent were added to the fresh sample and the AlCl₃ spectrum was recorded, 3 drops of HCl were added and after mixing, the AlCl₃/ HCl spectrum was recorded. Powdered NaOAc was then added to the fresh sample, the mixture was shaken and the NaOAc spectrum was recorded. NaOAc/ H₃BO₃ spectrum was then recorded after adding H₃BO₃.

Preparation of plant extract for phytochemical screening

Powdered shade-dried stem barks (200g) of *Detarium senegalense* were macerated with 95% ethanol for 48hr. The solvent

was removed *in vacuo* to give a crude extract used for phytochemical screening according to the method described by Harborne.

Extraction of flavonoids

Powdered shade –dried stem barks of *Detarium senegalense* (2 kg) were macerated with 95% ethanol at ambient temperature for 72 hours. The solvent was removed under reduced pressure to give a crude extract. The crude extract was dissolved in water and partitioned with ethyl acetate to afford the ethyl acetate fraction which was found to be rich in phenolics.

Paper chromatography (PC)

The ethyl acetate fraction of *Detarium senegalense* stem bark was dissolved in methanol and applied to Whatman paper (No. 3 mm – 46x 57cm). The bands were irrigated with BAW (n- butanol acetic acid-water, 5:2:6 ; v:v:v). The developed chromatograms were air-dried and examined under both visible and UV light (λ 366,245nm). The equivalent bands from each PC were then cut out, combined, cut into small strips and slurred with methanol. After several hours of contact with occasional shaking, the liquid was filtered and evaporated to dryness. In this way compound I was isolated in a chromatographically pure form.

Antimicrobial assay

Different fractions of *Detarium senegalense* stem bark were screened for their antimicrobial activity against four bacterial species, Gram-positive (*Staphylococcus aureus* and *Bacillus subtilis*), Gram-negative (*Pseudomonas aeruginosa* and *Escherichia coli*) and two fungal species (*Aspergillus niger*, *Candida albicans*). The cup plate agar diffusion bioassay was used.

Preparation of bacterial suspensions

One ml aliquots of a 24 hours broth culture of the test organisms were aseptically distributed on to nutrient agar slopes and incubated at 37° C for 24 hours. The bacterial growth was harvested and washed off with 100 ml sterile normal saline, to produce a suspension containing about 10⁸- 10⁹ C.F.U/ ml. The suspension was stored in the refrigerator at 4° C till used. The average number of viable organisms per ml of the stock suspension was determined by means of the surface viable counting technique.

Serial dilutions of the stock suspension were made in sterile normal saline solution and 0.02 ml volumes of the appropriate dilution were transferred by micro pipette onto the surface of dried nutrient agar plates. The plates were allowed to stand for two hours at room temperature for the drops to dry and then incubated at 37 °C for 24 hours. After incubation, the number of developed colonies in each drop was counted. The average number of colonies per drop (0.02 ml) was multiplied by 50 and by the dilution factor to give the viable count of the stock suspension, expressed as the number of colony forming units per ml suspension.

Each time a fresh stock suspension was prepared. All the above experimental conditions were maintained constant so that suspensions with very close viable counts would be obtained.

Preparation of fungal suspension

The fungal cultures were maintained on Sabouraud dextrose agar, incubated at 25 °C for 4 days. The fungal growth was harvested and washed with sterile normal saline and finally suspension in 100ml of sterile normal saline, and the suspension were stored in the refrigerator until used.

Testing of antibacterial susceptibility

The cup plate agar diffusion method was used to screen the antibacterial activity of plant extracts and performed by using Mueller Hinton agar (MHA). The experiment was carried out according to the National Committee for Clinical Laboratory Standards Guidelines (NCCLS, 1999). Bacterial suspension was diluted with sterile physiological solution to 10⁸cfu/ ml (turbidity = McFarland standard 0.5). One hundred microliters of bacterial suspension were swabbed uniformly on surface of MHA and the inoculum was allowed to dry for 5 minutes. Sterilized filter paper discs (Whatman No.1, 6 mm in diameter) were placed on the surface of the MHA and soaked with 20 μ l of a solution of each plant extract. The inoculated plates were incubated at 37 °C for 24 h in the inverted position. The diameters (mm) of the inhibition zones were measured [4].

Testing of antifungal susceptibility

The above-mentioned method was adopted for antifungal activity, but instead of agar, Sabouraud dextrose agar was used. Samples were used here by the same concentrations used above [5].

RESULTS

Phytochemical screening

Phytochemical screening of *Detarium senegalense* ethanolic extract revealed the presence of sterols, triterpenes, flavonoids, tannins, saponins, anthraquinones and cyanogenic glycosides. The ethyl acetate fraction was purified by paper chromatography. After the usual workup, compound I was isolated from *Detarium senegalense* stem bark in chromatographically pure form.

Identification of compound I

The IR spectrum of compound I (Fig.1) showed ν (KBr) 675 (CH, Ar. bending), 1023(CO), 1450, 1531(C=C, Ar.), 1645(C=O), 2923 (C-H, aliph.) and 2340cm⁻¹ (OH).

Compound I absorbs in the UV (Fig.2) at λ_{max} (MeOH) 280nm giving only band II. Such absorption- which originates from a benzoyl chromophore is revealed by: flavanones, isoflavones, dihydrchalcones and dihydroflavonols. These classes lack conjugation between the carbonyl function and the B ring. Thus only one band- band II is manifested. Other classes of flavonoids (flavones, flavonols, chalcones and aurones) reveal both bands I and II in their UV spectra due to conjugation between ring B and the carbonyl function at C-4. Such flavonoids possess two

chromophores: the benzoyl (giving band II) and cinnamoyl (giving band I) systems.

No shoulder in the range: 300-340nm was observed in the UV spectrum suggesting absence of isoflavones.

Next the hydroxylation pattern of this isolate was investigated using UV shift reagents: sodium methoxide, sodium acetate, aluminium chloride and boric acid. The ¹H NMR spectrum (Fig.7) showed δ(ppm):1.20(6H) assigned for two methyl functions, while the signal at 1.80(3H) was assigned for an acetyl group. The signal at δ3.73(3H) was attributed to a methoxyl function. The multiplet centered at δ3.50(6H) accounts for a sugar moiety. The signal at δ6.25(1H) accounts for C₆ – proton. Other aromatic protons appeared at δ6.73ppm [6], [7].

CONCLUSION

Detarium senegalense stem bark was screened for secondary metabolites. Paper chromatography allowed isolation of a dihydrochalcone from stem bark. On the basis of its spectral data, a partial structure was suggested for the isolate. Different fractions of *Detarium senegalense* stem bark were evaluated for antimicrobial activity against six standard human pathogens and significant results were obtained.

REFERENCES

1. Spångberg LSW, Haapasalo M, 2002. Rationale and efficacy of root canal medicaments and root filling materials with emphasis on treatment outcome. *Endod Top.* 2, Pages 35-58.
2. Wang Q, Ellis P.R., Ross-Murphy, S.B. and Burchard, W., *Carbohydrate Polymers.*, 1997, 31, 115-124. 12-Burkill, H.M., *Royal Bot. Gardens.* 3, Pages 102-105.
3. Gopikrishna AV, Kandaswamy D, Jeyaval Rajan K. 2006. Comparative evaluation of the antimicrobial efficacy of five endodontic root canal sealers against *Enterococcus faecalis* and *Candida albicans*. *J Cons Dent.* 9, Pages 2-11.
4. Manyahi J, Matee MI, Majigo M, 2014. Predominance of multi-drug resistant bacterial pathogens causing surgical site infections in Muhimbili National Hospital, Tanzania. *BMC Res Notes.* 7, Pages 500. Doi: 10.1186/1756-0500-7-500.
5. Muhammad UK, Adamu TM, Binji Z, 2014. Prevalence of β-lactamase production among pathogenic bacteria isolated from surgical site and wound infection among patients admitted in some selected hospitals in Sokoto Metropolis, Nigeria. *Int J Env.* 3(3), Pages 104-112.
6. Poirel L, Walsh TR, Cuvillier V, 2011. Multiplex PCR for detection of acquired carbapenemase genes. *Diagn Microbiol Infect Dis.* 70(1), Pages 119-123.
7. Sharif L, Obaidat M, Al-Dalalah M, 2013. Food Hygiene Knowledge, Attitudes and Practices of the Food Handlers in the Military Hospitals. *FNS.* Doi: <http://dx.doi.org/10.4236/fns.2013.43033>. J