



Research article

Antipyretic activity of ethanolic extract of Prosopis cineraria PODSAbhishek Pandey^{1, 2*}, Atul Kaushik², Suman Jain³, Bhagat Singh Jaiswal³¹ Department of Pharmaceutical Sciences, IFTM University, Moradabad, Uttar Pradesh, India² IPS College of Pharmacy, Gwalior, Madhya Pradesh, India³ School of Studies in Pharmaceutical Sciences, Jiwaji University, Gwalior, Madhya Pradesh, India**Corresponding author:** Abhishek Pandey, ✉

Department of Pharmaceutical Sciences, IFTM University, Moradabad, Uttar Pradesh, India

© The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by-nc/4.0/>). See <https://jmpas.com/reprints-and-permissions> for full terms and conditions.**Received** – 20 January 2016, **Revised** - 25 January 2016, **Accepted** – 23 February 2016 (DD-MM-YYYY)**Refer This Article**Abhishek Pandey, Atul Kaushik, Suman Jain, Bhagat Singh Jaiswal, 2016. Antipyretic activity of ethanolic extract of Prosopis cineraria PODS. Journal of medical pharmaceutical and allied sciences, V 6 - I 1, Pages -470 – 472. Doi: <https://doi.org/10.55522/jmpas.V6I1.0126>.**ABSTRACT**

Prosopis cineraria used in folk medicine to treat various ailments such as anemia, arthritis, asthma, toothache, ulcers and vertigo. Prosopis cineraria bark is used as remedy against the rheumatism, fever and inflammation. Pods is used for earache, toothache, pain relief from fractured bones. The aim of present study is to evaluate the antipyretic effect of Prosopis cineraria pods. Phytochemical screening of Prosopis cineraria pods were carried out to determine the presence of various phytoconstituents. Antipyretic effect was evaluated by Brewer's yeast induced pyrexia method. Phytochemical screening revealed presence of various secondary metabolites. Ethanolic extract of Prosopis cineraria pods (PCEE) show significant Antipyretic activity at a dose of 400 mg/kg. The results indicate that the ethanolic extract of Prosopis cineraria pods can be a good candidate for reducing fever and validate its use as a useful folk medicine due to the presence of various secondary metabolites.

Keywords: Pods, Pyrexia, Prostaglandins, Brewer's yeast, Arid zone.**INTRODUCTION**

Prosopis cineraria is locally known as "Khejri" in India. It is used in folk medicine to combat various ailments, viz., anaemia, arthritis, asthma, biliousness, boils, bronchitis, cholera, colic, toothache, ulcers and vertigo. It is reported that Prosopis cineraria bark is used traditionally as a remedy against rheumatism. Pods are consumed as food in the arid region of Rajasthan, additionally considered astringent. Aqueous paste of bark is applied externally to disinfect wounds and promote healing. Pharmacologically, it has demonstrated analgesic and anti-inflammatory activity. Fresh Leaves juice mixed with lemon juice is used for dyspepsia, the extract of crushed pods is used for earache, toothache, and pain relief from fractured bones. From a chemical point of view, the presence of phytosterols, flavonoids, tannins, phenols, carbohydrates, proteins and amino acids was detected in the preliminary phytochemical tests. Literature survey revealed that the antipyretic activity of Prosopis cineraria pods has not been scientifically proven. Therefore, the

present study aimed to evaluate the antipyretic activity of Prosopis cineraria pods [3].

MATERIALS AND METHODOLOGY**Chemicals and drugs**

Paracetamol (Plethico Pharmaceuticals, Indore) and brewer's yeast (HiMedia Laboratories Pvt Ltd, Mumbai, India), and other solvents were of analytical grade.

Collection and authentication of plant material

The fresh pods of Prosopis cineraria were collected from Jodhpur, Rajasthan, India and identified by a senior taxonomist. R.P. Pandey of National Research Institute for Ayurveda- Siddha Human Resource Development, Gwalior, India

Extraction and phytochemical screening

The pods of Prosopis cineraria were dried, chopped and ground by using a mechanical grinder to obtain a coarse powder and extracted with petroleum ether to remove fatty substances. Then finally extracted (cold maceration) with ethanol to obtain the ethanol extract. The extracts were filtered and concentrated over a water bath. The percentage yield of ethanolic extract was 6% w/w. Preliminary

phytochemical study of the extract was carried out to determine the presence of carbohydrates, proteins, flavonoids, alkaloids, terpenoids, fixed oils, and glycosides in the extract.

Animals

Healthy adult albino Wistar rats (200-225 g) of either sex were used for the study. The rats were acclimatised at a controlled room temperature $25 \pm 2^\circ\text{C}$ and 12 h light-dark cycle for 7 days. Food and water were given ad libitum. The protocol of the study was approved by the Institutional Animal Ethics Committee (IPS/COP/IAEC/01) according to the guidelines of the Committee for Control and Supervision of Experiments on Animals (CPCSEA), New Delhi, India.

Grouping and treatments

The animals were randomly divided into 4 groups 5 animals in each group. The PCEE were dissolved in 0.9% saline for oral administration.

Group I: Control group given 0.9% normal saline (p.o.).

Group II: Standard groups received (paracetamol 100 mg/kg, p.o.).

Group III & IV: PCEE (200 and 400 mg/kg, orally) treated groups.

Fever was induced by Brewer's yeast in normal saline 12.5% of 1 ml/100 gm body weight subcutaneously.

Antipyretic activity

Brewer's yeast induced pyrexia

Rats were fasted for 18 h before inducing pyrexia. The initial rectal temperature was noted by digital thermometer to 3 cm depth into the rectum of the animal. Brewer's yeast in normal saline 12.5% of 1 ml/100 gm body weight, subcutaneously administered to nap of the neck and the site of injection massaged to spread the suspension to induce fever. After 18 h the rise in rectal temperature was recorded. Animals show 1°C rise in body temperature were included in the study and administered with PCEE at dose (200 and

400 mg/kg) or paracetamol 100 mg/kg and 0.9% normal saline solution. The rectal temperature was recorded at 1, 2 and 3 h after dosing.

Statistical analysis

The results were analysed by two-way ANOVA and Bonferroni post hoc tests using Prism pad statistics software ver. 5. A statistical difference of $P < 0.05$ was considered significant in all test [4, 5].

RESULTS AND DISCUSSION

Phytochemical screening

Ethanol extract revealed the presence of carbohydrates, proteins, flavonoids, alkaloids, terpenoids, fixed oils, and glycosides in the extract.

Antipyretic Activity

PCEE in low dose 200 mg /kg, effective only at 3 hr ($p < 0.05$) to reduce the fever. While the PCEE dose of 400 mg/kg exhibited a significant antipyretic effect at 2 and 3 hr ($p < 0.01$, $p < 0.001$, respectively), a prominent effect of PCEE was observed at 3 hr. which is comparable to the standard drug paracetamol 100 mg/kg (Table 1). Brewer's yeast (lipopolysaccharide, which is the cell wall component of Gram-negative bacteria) is an exogenous pyrogen that binds to the immunological protein called the lipopolysaccharide binding protein. Fever induced by yeast is known as pathogenic fever. This binding results in the synthesis and release of various endogenous cytokine factors such as interleukin (IL) -1, IL-6, and TNF α , which activate the arachidonic acid pathway, and ultimately result in the synthesis and release of prostaglandin E2 (PGE2).[8-9] PGE2 is a key mediator of fever, acting on thermo-sensitive and thermo-integrative hypothalamic neurons. Inhibition of prostaglandin synthesis could be the possible mechanism of antipyretic action. Paracetamol act by inhibiting effect of pyrogen in preoptic region of the hypothalamus.

Table 1: Antipyretic effect of PCEE against Brewer's yeast induced pyrexia

Time (hrs)	Control group	Standard Paracetamol 100 mg/kg	PCEE 200 mg/kg	PCEE 400 mg/kg
0	98.48 \pm 0.19	98.10 \pm 0.32	98.88 \pm 0.58	98.71 \pm 0.2
18 h	101.31 \pm 0.25	101.70 \pm 0.51	101.34 \pm 0.42	101.38 \pm 0.30
1	101.44 \pm 0.48	99.81 \pm 0.28**	101.56 \pm 0.55	100.9 \pm 0.26
2	101.66 \pm 0.22	98.29 \pm 0.60***	100.88 \pm 0.45	99.88 \pm 0.57**
3	101.38 \pm 0.18	98.68 \pm 0.39***	100.12 \pm 0.29*	99.31 \pm 0.37***

Two way ANOVA Non repeated measure using Bon Ferroni post hoc test * $p < 0.05$ ** $p < 0.01$ *** $p < 0$.

PCEE produce antipyretic effects in higher dose 400 mg/kg comparable to the paracetamol treated group which may be due to inhibition of prostaglandin synthesis.

Phytochemicals present in pods ethanolic extract such as flavonoids, alkaloids, terpenoids, fixed oils, glycosides may be responsible for antipyretic effect [6].

CONCLUSION

In conclusion, Prosopis cineraria pods enriched with antipyretic activity and affirm its use as a remedy for the treatment of fever. However, the isolation of compounds responsible for antipyretic

effect is future scope of the study.

Declaration of interest

The authors declare that there is no conflict of interest.

ACKNOWLEDGMENT

The authors acknowledge director SOS in pharmaceutical sciences, Jiwaji University, Gwalior for their encouragement and technical support.

REFERENCES

1. Akash Garg, 2013. Review on Prosopis cineraria: A potential herb of Thar desert. Drug invention today. 5, Pages 60-65. Doi: <https://doi.org/10.1016/j.dit.2013.03.002>.

2. Nidhi Sharma, 2010. Anti hyperglycemic antihyperlipidemic and antioxidative potential of *Prosopis cineraria* bark. *Indian Journal of Clinical Biochemistry*. 2 (2), Pages 193-200.
3. Sumitra Singh, 2013. Pharmacognostical and phytochemical studies on stem bark of *Prosopis cineraria*- A medicinal plant indigenous to South West Asia. *Universal Journal of Pharmacy*. (02), Pages 173-180.
4. Khandelwal KR, Practical Pharmacognosy, 2006. 15th ed. Pune: Nirali Prakashan. Pages 149-156
5. Ukwuani AN, Abubakar MG, Warra SH, 2012. Antipyretic activity of some Nigerian medicinal plants in rats. *International Journal of Pharmaceutical Research*. 4, Pages 48– 51.
6. Aman AA, Patrick NO, 2011. Investigation of anti-inflammatory, antipyretic and analgesic effect of Yemeni Sidr honey. *World Academy of Science Engineering and Technology*. 80. Pages 47– 52.
7. Ashok BK, Ravishankar B, Prajapati PK, 2010. Antipyretic activity of Guduchi Ghrita formulations in Albino Rats. *Ayu*. 31, Pages 367–3.
8. Li S, Dou W, Tang Y, 2008. Acetaminophen: antipyretic or hypothermic in mice? In either case, PGHS-1b (COX-3) is irrelevant. *Prostaglandins Other Lipid Mediators*. 85, Pages 89–99. Doi: 10.1016/j.prostaglandins.2007.10.007
9. Abdul Qadir Panhwar, 2007. Ethnobotanical studies of Mahal Kohistan (Khirthar National Park) Pakistan *Journal of Botany*. 39(7), Pages 2301-2315.